Skeletal muscle mitochondrial function cannot be properly inferred from PCr resynthesis without taking pH changes into account

Sebastien Ratel, Vincent Martin, Anne Tonson, Patrick J. Cozzzone, David Bendahan

► To cite this version:

Sebastien Ratel, Vincent Martin, Anne Tonson, Patrick J. Cozzzone, David Bendahan. Skeletal muscle mitochondrial function cannot be properly inferred from PCr resynthesis without taking pH changes into account. *Magnetic Resonance Imaging*, 2012, 30 (10), pp.1542 - 1543. 10.1016/j.mri.2012.06.017 . hal-01659908

HAL Id: hal-01659908

<https://hal.uca.fr/hal-01659908>

Submitted on 9 Dec 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ELSEVIER

Available online at www.sciencedirect.com

SciVerse ScienceDirect

Magnetic Resonance Imaging xx (2012) xxx–xxx

MAGNETIC
RESONANCE
IMAGING

Skeletal muscle mitochondrial function cannot be properly inferred from PCr resynthesis without taking pH changes into account

To the Editor:

We have read with much interest the paper recently published by Khushu et al. regarding the impaired muscle metabolism in hypothyroid patients [1]. Although the results of this investigation are of potential interest, we are actually concerned about the utilization of the phosphocreatine recovery rate constant (kPCr) as a simple index of mitochondrial function and the corresponding conclusions regarding the impact of hypothyroidism on muscle metabolism during exercise. The real issue is actually to clearly determine whether one can infer about mitochondrial function on the basis of PCr rate constants measurements.

During recovery from exercise, PCr is resynthesized purely as a consequence of oxidative adenosine triphosphate (ATP) synthesis [2], and measurements of the kPCr have been used in order to characterize mitochondrial function in a variety of conditions [3]. However, several studies have clearly demonstrated that cytosolic pH has a strong influence [3]. It is actually well acknowledged that kPCr is inversely related to the extent of intracellular acidosis and PCr consumed [3]. In other words, a significant intracellular acidosis and a large PCr consumption in exercising muscle would be associated with a slower PCr resynthesis [3]. In the Khushu et al. study, cytosolic pH was found to be significantly lower after a 6-min constant-load plantar flexion exercise in hypothyroid patients compared to controls (6.95 ± 0.02 vs. 7.02 ± 0.06 , respectively). Therefore, the conclusions of an altered mitochondrial function in hypothyroid patients [1] on the single basis of kPCr measurements have to be interpreted cautiously. Of note, several kinetic parameters are commonly used to describe PCr changes during exercise-to-recovery transition, including kPCr, the initial rate of PCr recovery (ViPCr) and the maximum aerobic capacity (Qmax) [4]. These three parameters characterizing PCr resynthesis are correlated to oxidative capacity. However, in contrast to kPCr, ViPCr and Qmax are insensitive to exercise intensity and end-of-exercise metabolic conditions [3]. On that basis, ViPCr and Qmax should be considered as additional indices to compare the postexercise PCr recovery rate and mitochondrial

oxidative capacity across different populations when end-of-exercise pH and PCr concentration values are different or not taken into account. These additional parameters were considered in some previously published studies which investigated in vivo mitochondrial function from childhood to young and/or late adulthood [5,6]. In contrast, these parameters were not taken into account in other studies which investigated age-related changes in mitochondrial oxidative capacity from childhood to adulthood [7] and examined the association between mitochondrial alterations and insulin sensitivity in overweight and normal-weight children [8]. In the Khushu et al. study, the Qmax calculation was oversimplified as the product of kPCr and the resting [PCr], as previously described [9]. This oversimplification is actually derived from the calculation of the ViPCr from an exponential fitting. In that case, ViPCr is represented by the product of kPCr and the amount of PCr consumed at the end of exercise. On that basis, a few authors have calculated Qmax similarly to ViPCr, with the amount of PCr consumed being replaced by the resting [PCr]. Although mathematically plausible, this calculation is not supported by any control model, and the corresponding Qmax is devoid of any enzymatic meaning, as it is the case for the calculation of Qmax using ADP within a Michaelis–Menten framework. This method is then largely questionable because it ignores the commonly accepted theory considering adenosine diphosphate as a key regulator of oxidative ATP synthesis [10]. In addition, with such an approximate calculation, Qmax becomes strongly dependent on kPCr and/or [PCr] measured at rest. In their study, Khushu et al. [1] reported no significant difference in the resting [PCr] values between hypothyroid patients and controls (43.6 ± 0.8 vs. 42.0 ± 1.0 mmol kg⁻¹, respectively). Given that the resting [PCr] values were similar, the corresponding calculated Qmax values were lower in hypothyroid patients than in controls on the single basis of lower kPCr measurements in hypothyroid patients. Consequently, this result should be interpreted with caution.

Overall, it is of utmost importance to keep in mind that one cannot simply infer on mitochondrial function on the basis of measurements of the rate constant of PCr resynthesis given that end-of-exercise conditions have been shown to exert a strong influence. Future studies should then pay attention to these methodological inaccuracies which might hedge data interpretation and confound the corresponding conclusions.

Sébastien Ratel

Vincent Martin

Clermont Université, Université Blaise Pascal, EA 3533
 Laboratoire des Adaptations Métaboliques à l'Exercice
 en conditions Physiologiques et Pathologiques (AME2P)
 BP 80026, F-63171 Aubière Cedex, France
 E-mail address: Sebastien.RATEL@univ-bpclermont.fr

Anne Tonson

Patrick J. Cozzone

David Bendahan

Centre de Résonance Magnétique Biologique et Médicale
 (CRMBM), Faculté de Médecine de Marseille, Université
 de la Méditerranée, Marseille, France

References

- [1] Khushu S, Rana P, Sekhri T, Sripathy G, Tripathi RP. Bio-energetic impairment in human calf muscle in thyroid disorders: a 31P MRS study. *Magn Reson Imaging* 2010;28:683–9.
- [2] Quistorff B, Johansen L, Sahlin K. Absence of phosphocreatine resynthesis in human calf muscle during ischaemic recovery. *Biochem J* 1993;291:681–6.
- [3] Roussel M, Bendahan D, Mattei JP, Le Fur Y, Cozzone PJ. 31P magnetic resonance spectroscopy study of phosphocreatine recovery kinetics in skeletal muscle: the issue of intersubject variability. *Biochim Biophys Acta* 2000;1457:18–26.
- [4] Tonson A, Ratel S, Le Fur Y, Vilmen C, Cozzone PJ, Bendahan D. Muscle energetics changes throughout maturation: a quantitative 31P-MRS analysis. *J Appl Physiol* 2010;109:1769–78.
- [5] Barker AR, Welsman JR, Fulford J, Welford D, Armstrong N. Muscle phosphocreatine kinetics in children and adults at the onset and offset of moderate-intensity exercise. *J Appl Physiol* 2008;105:446–56.
- [6] Taylor DJ, Kemp GJ, Thompson CH, Radda GK. Ageing: effects on oxidative function of skeletal muscle in vivo. *Mol Cell Biochem* 1997;174:321–4.
- [7] Fleischman A, Makimura H, Stanley TL, et al. Skeletal muscle phosphocreatine recovery after submaximal exercise in children and young and middle-aged adults. *J Clin Endocrinol Metab* 2010;95:E69–74.
- [8] Fleischman A, Kron M, Systrom DM, Hrovat M, Grinspoon SK. Mitochondrial function and insulin resistance in overweight and normal-weight children. *J Clin Endocrinol Metab* 2009;94:4923–30.
- [9] Conley KE, Jubrias SA, Esselman PC. Oxidative capacity and ageing in human muscle. *J Physiol* 2000;526:203–10.
- [10] Kemp GJ, Taylor DJ, Radda GK. Control of phosphocreatine resynthesis during recovery from exercise in human skeletal muscle. *NMR Biomed* 1993;6:66–72.

AUTHOR QUERY FORM

ELSEVIER	Journal: MRI Article Number: 7852	Please e-mail or fax your responses and any corrections to: Nina Biuso E-mail: nbiuso@aol.com Tel: 845-896-0870 Fax: 845-896-0870
---	--	--

Dear Author,

Please check your proof carefully and mark all corrections at the appropriate place in the proof (e.g., by using on-screen annotation in the PDF file) or compile them in a separate list. Note: if you opt to annotate the file with software other than Adobe Reader then please also highlight the appropriate place in the PDF file. To ensure fast publication of your paper please return your corrections within 48 hours.

For correction or revision of any artwork, please consult <http://www.elsevier.com/artworkinstructions>.

Any queries or remarks that have arisen during the processing of your manuscript are listed below and highlighted by flags in the proof. Click on the 'Q' link to go to the location in the proof.

Location in article	Query / Remark: click on the Q link to go Please insert your reply or correction at the corresponding line in the proof
Q1	Please provide postal code here. <div data-bbox="540 1016 1065 1134" style="border: 1px solid black; padding: 5px; margin: 10px auto; width: fit-content;"><p>Please check this box if you have no corrections to make to the PDF file. ☐</p></div>

Thank you for your assistance.