1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Validation of a new scoring scale for behavioral assessment of L-dopa-induced dyskinesia in the rat: a new tool for early decision-making in drug development

Simon Loiodice^{1,*}, Anne-Sophie Denibaud¹, Wendy Deffains¹, Magali Alix¹, Pierre Montagne¹,
Marine Seffals², Christophe Drieu La Rochelle¹

¹Non-Clinical Department, Biotrial Pharmacology, 7-9 rue Jean-Louis Bertrand 35042 Rennes, France

²Plate-Forme H2P2, Université de Rennes 1, Biosit, 2 Av. du Prof. Léon Bernard, 35043 Rennes, France.

*Corresponding Author:

Simon Loiodice

Non-Clinical Department
Biotrial Pharmacology
7-9 rue Jean-Louis Bertrand
35042 Rennes, France
Email: simon.loiodice@biotrial.com

Abstract

The 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP)-treated non-human primate (NHP) has been described as the most translatable model for experimental reproduction of L-dopa-induced dyskinesia (LID). However, from a drug discovery perspective, the risk associated with investment in this type of model is high due to the time and cost. The 6-hydroxydopamine (6-OHDA) rat dyskinesia model is recommended for testing compounds but relies on onerous, and non-standard behavioral rating scales. We sought to develop a simplified and sensitive method aiming at assessing LID in the rat. The purpose was to validate a reliable tool providing earlier insight into the antidyskinetic potential of compounds in a time/cost-effective manner before further investigation in NHP models.

Unilaterally 6-OHDA-lesioned rats were administered L-dopa (20 mg/kg) and benserazide (5 mg/kg) daily for 3 weeks starting 4 weeks post-lesion, then co-administered with amantadine (20-30-40 mg/kg). An adapted rating scale was used to score LID frequency and a severity coefficient was applied depending on the features of the observed behavior.

A gradual increase (about 3-fold) in LID score was observed over the 3 weeks of L-dopa treatment. The rating scale was sensitive enough to highlight a dose-dependent amantadine-mediated decrease (about 2.2-fold) in LID score.

We validated a simplified method, able to reflect different levels of severity in the assessment of LID and, thus, provide a reliable tool for drug discovery.

Keywords: Parkinson's disease; L-dopa; Dyskinesia; 6-OHDA; Drug discovery

Introduction

To date, L-dopa is the most effective drug for treating the signs and symptoms of Parkinson’s disease (PD) and is still considered the standard therapeutic agent ^{1,2}. However, after several years of treatment, L-dopa-induced dyskinesia (LID) occurs in the majority of patients ^{1,3–5}. These highly debilitating motor complications manifest as a variety of abnormal involuntary movements (AIMs), including severe and painful dystonic cramps (sustained abnormal muscle contractions), hyperkinetic and purposeless (choreiform) movements ^{5–7}.

The pathophysiological mechanisms underlying LID remain poorly understood despite substantial efforts in research ^{8,9}. LID has been proposed to be an aberrant form of neuroplasticity triggered by the combined effects of dopaminergic (DA) denervation and pulsatile stimulation of the DA receptors through repeated L-dopa intake ^{1,6}. With increasing duration of treatment, there is an increase in both the frequency and the severity of dyskinesia ⁶. Given that L-dopa is still required for the treatment of advanced PD, the management of LID is a crucial challenge ^{8,9}. The limited success of anti-dyskinetic drugs in the clinic emphasizes the need to use appropriate animal models with good predictive value. The MPTP-treated non-human primate (NHP) has been described as the most translatable model and the gold standard for experimental reproduction of PD features, as well as dyskinetic syndrome ^{7,10,11}. However, the use of NHPs is costly, time-consuming and raises potential ethical issues. Hence, a drug candidate failing due to lack of efficacy in NHP dyskinesia models may result in losing a significant investment in the research program (potentially resulting in significant consequences for a small biotech company, and in no new drug being available for patients). In agreement with this idea, it has been proposed that early attrition of non-efficacious compounds is crucial for successful central nervous system (CNS) drug development ^{12,13}.

To counteract the limitations associated with the use of NHPs in antidyskinetic drug discovery, other simpler models were developed. The unilateral 6-hydroxydopamine (6-OHDA) rat dyskinesia model remains one of the most popular for modeling peak-dose dyskinesia ⁷. This model was first described

by Cenci and colleagues in 1998¹⁴ and is still widely used and recommended for testing compounds, because it produces a stable and reproducible behavioral outcome^{7,15}. This model relies on a behavioral assessment of LID using a rating scale that includes measurement of locomotor behavior (contralateral turning) as well as abnormal involuntary movements (AIMs), such as forelimb dyskinesia, dystonic posturing of the contralateral side of the body, axial dystonia, manifest as contralateral twisted posture of the neck and upper body, and orolingual dyskinesia, characterized by stereotyped jaw movements and contralateral tongue protrusion¹⁴. Lundblad and colleagues elegantly demonstrated progressive worsening of AIMs over a 3-week L-dopa treatment using this scoring method¹⁶. However, the severity of the AIMs was determined based on the frequency of occurrence of different subtypes of behaviors, without taking into account their intensity level, which is known to increase over time along with LID frequency with continued L-dopa treatment in PD patients, as reviewed by Bezard and colleagues⁶. This limitation was later counterbalanced by adding an amplitude score (based on the extent of the movements) in order to differentiate between small but clear forelimb movements and dystonic-like movements involving the whole shoulder^{17,18}. Another rating scale aimed at further assessing the intensity of LID was meanwhile developed by Steece-Collier and colleagues¹⁹. Interestingly, in this latter method, the score was assigned not only based on the duration of a behavior, but also on its features, in order to better reflect LID severity¹⁹. All of these rating scales have recently been compared in a validation study, in order to assess their reliability and translatability²⁰. The authors obtained slight differences in the responses to reference compounds depending on the rating scale, and suggest (as do other authors) a need to standardize the method^{7,20}. Notably, they stated that methods enabling the detection of variations in both duration and intensity of LID resulted in improved sensitivity to the effect of the reference compound amantadine, a non-competitive NMDA receptor antagonist⁷.

In clinical research, it has been highlighted that the development of a single sensitive and robust rating scale was challenging because of the different types of dyskinesias and their different temporal patterns, anatomical distributions and associated disabilities⁸. Similarly, although well

validated, the different methods described above reflect a lack of harmonization in the behavioral assessment of LID in animal models. Most of the existing scales are relevant for profound investigations into the neurobiological mechanisms of LID (if not designed specifically for this purpose) and rely on direct visual observation of a large number of fine behavioral parameters. As a consequence, the impact of subjectivity is high (the experimenter may introduce a subjective bias for each behavioral subtype observed, which influences the global LID score), and the use of a particular method can be challenging for an experimenter who was not trained by the laboratory in which the method was developed. This suggests a need to develop an 'easy-to-use' LID scoring scale that could be used in the 6-OHDA rat as a de-risking strategy, to have earlier insight into drug efficacy before carrying out further expensive studies using NHP models of dyskinesia.

In the present study, we sought to develop a new scoring method, adapted from previous rating scales^{14,16,17,19–22} for the assessment of LID in the rat 6-OHDA dyskinesia model. The purpose was to validate a simplified method focusing on a reduced number of behavioral subtypes easily identified in order to lower the impact of subjectivity and provide reliable data concerning the antidyskinetic potential of a compound in a time/cost-effective manner.

RESULTS & DISCUSSION

As shown in **Fig. 2a** and **Fig. 2b**, the animals that received a stereotaxic injection of 6-OHDA showed a significant decrease in the number of TH+ cells within the SNc (81.1% cell loss compared to the unlesioned side, $t_{34}=11.09$; $p<0.0001$). However, 2 rats showed a more partial lesion, of 31% and 44%, respectively (black dots in **Fig. 2b**). TH immunostaining performed in 2 sham animals revealed a number of 22,267 and 15,800 TH+ cells in the SNc, respectively, which was similar to the numbers of TH+ cells observed in the unlesioned side of lesioned rats (18,696 on average). Forelimb use asymmetry and akinesia were assessed using the cylinder test and the stepping test, respectively. As

shown in **Fig. 2c**, the percentage of contralateral forelimb paw use was significantly decreased in 6-OHDA-lesioned rats (29.4%) compared to sham rats (48.4%) ($p=0.0005$). This motor impairment was reversed after treatment with L-dopa (8 mg/kg) co-administered with benserazide (6 mg/kg). Consistently, the stepping test procedure reflected a similar motor impairment that was alleviated with the same L-dopa treatment (**Fig. 2d**). Indeed, while sham rats made an average of 23.8 adjusting steps with their contralateral paw over 3 trials, vehicle-treated lesioned rats made only 4.4 steps ($p<0.0001$) and L-dopa-treated lesioned rats made 16.8 steps ($p<0.0001$ compared to vehicle, $p=0.0057$ compared to sham). **Fig. 2e**, reveals that apomorphine challenge significantly induced contralateral rotations, especially in 8 rats, for which the challenge induced over 90 turns/45 min. Based on immunohistochemistry analysis, these rats underwent a dopaminergic lesion causing a TH+ cell loss of over 80% compared to the unlesioned side and were selected for the assessment of L-dopa-induced dyskinesia (apomorphine did not produce contraversive turn in 10 rats including 2 partially lesioned rats represented by black dots in **Fig. 2e** and **Fig. 2b**).

Drug-induced dyskinesia is a frequent debilitating complication in PD, associated with physical and social disabilities. For decades, substantial research efforts have been invested into novel therapeutics that could improve patients' quality of life, with limited success. Consistent with the need to develop standardized and cost/time-effective preclinical tools for anti-dyskinetic drug discovery, we provide evidence highlighting the value of a novel 'easy-to-use' reliable method for behavioral assessment of LID in the rat.

The late stage (Phase II, Phase III) attrition rate of CNS drug-candidates is known to be particularly high, mainly due to lack of efficacy (or safety issues)^{12,13,23,24}. The successful development of CNS drugs relies on the development of appropriate animal models for efficacy testing in preclinical studies and a new paradigm for drug development that will give early readouts for proof of concept in order to allow attrition to occur much earlier in the process^{12,13}. This challenge applies to PD drug

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

discovery, especially for the development of antidyskinetic drugs ⁷. To date, the MPTP-treated NHP is considered the most predictive animal model of LID ^{7,10,11}. Indeed, after chronic treatment with L-dopa, MPTP-treated monkeys develop choreic and dystonic movements that can be scored using clinical rating scales after only modest adaptations ²⁵. But the cost and time required to implement this type of studies is associated with a risk of losing a significant investment. The 6-OHDA dyskinesia rat model appears to be a good option to provide a first insight into drug efficacy in a time/cost-effective manner ¹¹. This model has been extensively described as valuable for antidyskinetic drug discovery ^{7,11,20} and relies on behavioral scoring of the LID. A recent study sought to validate 3 contemporary AIM scales available in the literature ²⁰ and highlighted the interest of adding a severity parameter to the scoring of AIM frequency. The scoring scale developed in our study attempts to take this crucial parameter into account. Both severity and adjustment coefficients were applied to a set of specific behavioral subtypes that were selected to reflect different levels of severity (**Table 1**). The goal was to provide an overview of the dyskinetic state of each animal rather than a very deep behavioral assessment. However, we designed our scale to reflect as much contrast as possible within the different levels of AIM severity to increase its sensitivity. It is worth noting that such an approach, relying on assigning a severity coefficient depending on the features of the observed behavior has been used in previous validated scales ¹⁹.

The evaluation of the evolution of the LID over the 3-week period of L-dopa treatment revealed a significant time-dependent increase in the LID score from week 0 to week 3 ($p=0.0362$) (**Fig. 3a**). It is worth noting that the LID scores correspond to ranked data. Hence, a square transformation was performed in order to ensure normal distribution and apply parametric tests. **Fig 3b** shows that “increased locomotion with contralateral bias” and “dystonia” were the main AIM components of the global LID score at week 0 and week 3 respectively (score significantly higher compared to other components). Furthermore, **Fig 3b** illustrates a significant increase of the score for “increased locomotion with grabbing” on weeks 1 and 2 ($p=0.0008$, $p=0.0038$) compared to week 0. On week 3, a significant increase of the score for “axial posture” ($p=0.0009$) and “dystonia” ($p=0.0151$) was

observed compared to week 0. **Fig. 3c** shows a dose-dependent decrease in the LID score when the animals received amantadine, with a maximum effect at the 40 mg/kg dose ($p=0.0034$). **Fig. 3d** illustrates that at this dose, most of the effect occurred between 30 min and 70 min post L-dopa/benserazide treatment ($p=0.007$ at 30 min, $p=0.0019$ at 40 min; $p=0.002$ at 50 min and $p=0.0155$ at 70 min after L-dopa/benserazide administration). It is worth noting that the 30 mg/kg dose produced a significant decrease in the LID score at 50 min ($p=0.0443$). **Fig. 3e** further illustrates the effect of the active dose of amantadine (40 mg/kg) on individual LID scores. **Fig. 3f** shows the contribution of the different behavioral subtypes of the scale to the global LID score among the different treatment groups after the 3-week period of L-dopa treatment. It reveals that “dystonia” and “axial posture” constituted the main components of the global LID score and the main parameters that were alleviated by amantadine since a significant decrease was observed for “dystonia” score at 30 mg/kg ($p=0.0379$) and 40 mg/kg ($p=0.0350$). It worth noting that the “axial posture” score was not alleviated at 30 mg/kg and slightly, but non-significantly, improved at 40 mg/kg.

Our data demonstrate that our rating scale was able to distinguish slight behavioral changes, as shown by the measurement of the time-dependent worsening of LID. Interestingly, we reported a significant increase of the LID score after one week of L-dopa dosing, consistent with a previous report indicating that AIM and LID can occur after the first-ever dose of L-dopa²⁶. Furthermore, the analysis of the different behavioral subtypes separately suggests a moderate increase of the most severe AIM components (i.e. dystonic circling, axial posture and dystonia) on weeks 1 and 2 accompanied by a significant augmentation of contralateral turning with grabbing. The two AIM subtypes “axial posture” and “dystonia” were significantly increased on the week 3 post-L-dopa treatment. This further illustrates the capability of our rating scale to highlight a progressive worsening of the dyskinetic state suggesting the sensitivity of the method. Moreover, the sensitivity of our rating scale allowed to show the amantadine-mediated alleviation of AIMS in a dose-dependent manner (using a tight dose range). It worth noting that amantadine-mediated alleviation

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

of the global LID score was mainly associated with an improvement of “dystonia” and, in a lesser extent, “axial posture” AIM components with very limited effect on other behavioral subtypes observed in our rating scale. This is in line with previous data reporting amantadine-mediated alleviation of AIM score with no effect of contralateral turning²⁰. Consistent with previous reports²⁰, our study illustrates the importance of performing assessments at multiple time-points, as we were able to pinpoint the time window corresponding to the optimal effect of amantadine. This is particularly relevant in experiments investigating compounds for which poor pharmacokinetics data are available, as some agents may reduce the severity of the AIM score at a peak time but prolong the temporal course of motor dysfunction, or have a delayed effect²⁰.

One of the reasons proposed to explain the limited success of anti-dyskinetic drugs in the clinic is the lack of validated clinical outcome measures that are responsive to treatment despite the validity of multiple dyskinesia rating scales²⁷, and it has been pointed out that the development of a single sensitive and robust rating scale was challenging because of the different types of dyskinesias and their different temporal patterns, anatomical distributions and associated disabilities⁸. Furthermore, the clinical assessment of LID is highly subject to the placebo effect²⁸. Similarly, in rat models of dyskinesia, various rating scales were developed and validated over the last decades^{14,16–19,21}. These methods rely on direct visual observation of a limited number of behavioral subtypes including orolingual/jaw movements (easily impacted by subjectivity) but only taking into account the severity level of AIM through frequency scores¹⁴ or amplitude scores depending on the observation of parameters easily subjectively appreciated (*e.g.* angle of torsion in moving animal)¹⁸. Other scales using a simpler method to determine a severity score focus on a large number of various fine subtypes of behavior or on observation of different parts of the animal’s body separately (neck, trunk, limbs)¹⁹. While these methods are appropriate for investigating the neurobiological mechanisms of LID their subjectivity remains high: a large number of subjective parameters may increase the chance of experimenter-dependent errors in assessment influencing an animal’s global dyskinetic score. Similarly, assessment of very fine behavioral parameters such as orolingual

movements or torsion angle in moving rats may introduce a subjective bias, impact the reproducibility and harm to the need of standardization. The reproducibility could only be ensured by significant training provided by the laboratory in which the method was first validated, as well as internal validation in the laboratory where the new research will be carried out. In a recent study comparing different rating scales, the authors suggested a need for standardization of the methods for assessing LID, as they obtained slight differences in the responses to reference compounds depending on the method used ²⁰. We believe that focusing on a limited number of behavioral subtypes could lower the impact of subjectivity and improve reproducibility while providing a global overview of the dyskinetic state of each animal. In our study, we observed a restricted number of behavioral subtypes to (i) minimize the impact of subjectivity (assuming that, as subjective bias would be applied to a smaller number of behavioral subtypes, there would be a lower impact on the final LID score due to fewer chances to produce “error”) and to (ii) obtain an overall view of the dyskinetic state of the animal at a particular time-point. Orolingual AIM were considered as too easily impacted by subjectivity (potentially hardly identified in dyskinetic moving rats) and were not assessed to minimize the risk to introduce a subjective bias. Our data demonstrate that this parameter may not be required to have a first insight into the antidyskinetic potential of a drug. We believe that this type of method could be more easily harmonized between different laboratories and meet expectations for the preclinical assessment of developing compound. Indeed, the purpose of our scale is not to support mechanistic studies aiming at understanding the neurobiological mechanisms of LID, but rather to provide, from a drug discovery perspective, a valuable, time/cost-effective outcome. While different subtypes of AIM may be mediated by different neurobiological mechanisms, assessing the effect of a test compound on the “global dyskinetic state” may help as part of a de-risking strategy to support decision-making during drug development (e.g., a go/no-go decision for initiating NHP studies). Our rating scale focuses on a limited number of behavioral subtypes, including non-dyskinetic behavior (contralateral circling) as well as AIMs with dystonic and hyperkinetic features. The behavioral subtypes assessed were associated with severity coefficients in

order to reflect the effect of a test compound on the global dyskinetic state of an animal. Our data demonstrate a time-dependent worsening of LID and a dose-dependent alleviation by amantadine, illustrating the sensitivity of this method.

In the present study, we provided data supporting the reliability of our new LID rating scale. However, as for other scales, the behavioral assessment relies on subjective observation of the animal, which is the main limitation of this type of a tool. The development of methods relying on more objective readouts, such as fluidic biomarkers or EEG monitoring, would be of major interest. For instance, substantial efforts have been made to identify gene/protein expression changes associated with LID in 6-OHDA animal models of PD^{29–31}, while other studies have attempted to identify fluidic biomarkers in PD patients³². Other EEG-based approaches have pointed out different specific patterns that are potentially associated with LID in humans^{33–35} and in the 6-OHDA rat model^{36,37}. It could be of great interest to combine similar approaches with our behavioral rating scale and assess whether results obtained using these methods correlate with each other and provide valuable information during the assessment of drug efficacy.

Many companies are trying to reduce costs by outsourcing drug discovery to academic labs or contract research organizations due to unfavorable risk/reward balance with CNS drug development^{23,38–40,41,41}. These new partners need to develop standardized and accessible tools for CNS drug discovery. Increasing confidence in preclinical data during the development of anti-dyskinetic agents is of crucial importance. In the present study, we provide data validating a simplified method that is able to reflect different levels of severity when assessing LID. The sensitivity of the rating scale allowed us to highlight a time-dependent worsening of LID during 3 weeks of L-dopa treatment and a dose-dependent reversion of LID by amantadine. We propose this method as a valuable, reliable and sensitive tool for the initial testing of novel antidyskinetic compounds. Although the method proposed here may not be appropriate for investigations on the neurobiological mechanisms of LID (no detailed assessment of various AIM subtypes), it may provide early efficacy data to support

1
2
3 decision-making during the drug development process. Indeed, early attrition of inefficacious
4
5 compounds is thought to be crucial for successful drug development (especially for CNS drugs),
6
7 hence the need to develop reliable methods that can rapidly provide efficacy data in a cost-effective
8
9 manner, before moving the compound forward in expensive and time-consuming studies. We believe
10
11 that the use of our rating scale in the 6-OHDA dyskinesia model can be valuable if integrated in a de-
12
13 risking strategy during the preclinical development of an antidyskinetic agent.
14
15

16 **Materials & Methods**

17 **Animals**

18
19
20 Adult male Sprague-Dawley rats (175-200 g) from Janvier Labs (Saint Berthevin, France) were
21
22 maintained in a controlled environment (lights on 07:00-19:00, $\pm 22^{\circ}\text{C}$) with food and water freely
23
24 available. They were housed 3-4 per cage. This study was carried out in AAALAC-accredited facilities
25
26 in strict accordance with the European Communities Council Directive (2010/63/EU) guidelines for
27
28 the care of laboratory animals. The protocol was approved by the Biotrial Pharmacology Committee
29
30 on the Ethics of Animal Experiments "Comité de Réflexion Ethique en Expérimentation Animale"
31
32 (CR2EA), and in accordance with French Research Ministry regulations. All possible efforts were
33
34 made to minimize suffering.
35
36
37
38
39
40

41 **Unilateral 6-OHDA lesion**

42
43
44 Twenty-six (26) rats were anesthetized with xylazine (Rompun®) 10 mg/kg, i.p., and ketamine
45
46 (Imalgen®) 80 mg/kg, i.p., before being placed in a stereotaxic frame (David Kopf Instruments, CA,
47
48 USA). The animals received a stereotaxic injection of 6-OHDA (4 $\mu\text{g}/\mu\text{L}$) or vehicle into the medial
49
50 forebrain bundle. This injection consisted of a 2.5 μL deposit at the following coordinates (from
51
52 bregma): AP: -4.4 mm; ML: -1.8 mm; DV: -7.9 mm (tooth bar at -2.4 mm) according to the Paxinos rat
53
54
55
56
57
58
59
60

1
2
3 brain atlas ⁴². 6-OHDA or vehicle was administered at a rate of 0.4 µL/minute. To avoid reflux, the
4
5 needle was maintained at the injection site for 5 minutes after the injection.
6

7
8 To limit damage to noradrenergic neurons, imipramine (15 mg/kg, i.p.) was administered 15 min
9
10 before 6-OHDA lesioning ⁴³.
11

12
13 The extent of the lesion was evaluated 3 weeks post-surgery, based on net apomorphine-induced
14
15 contralateral rotations ^{43,44}. Animals were injected with apomorphine (0.05 mg/kg, s.c.), immediately
16
17 placed in a 45 x 45 cm plexiglas open-field and video-tracked over 45 min. The number of
18
19 contralateral rotations was measured afterwards using EthoVision® XT 9.0 (Noldus, Netherlands).
20
21 Animals with a number of rotations greater than 90 were considered to have greater than 95% DA
22
23 cell loss ^{43,44}.
24

25
26 Prior to the cylinder and stepping tests (see *Behavioral procedures*), lesioned animals were
27
28 randomized according to their number of contralateral turns in order to have a homogeneous
29
30 distribution between the L-dopa- and saline-treated groups.
31

32
33 **Drugs**
34
35

36
37 All drugs (Sigma, France) were dissolved in saline, except 6-OHDA, which was dissolved in a saline
38
39 solution containing 0.02% ascorbic acid.
40

41
42 In the cylinder and stepping tests, L-dopa methyl ester and benserazide were administered at 8 and
43
44 6 mg/kg, ip, respectively ^{45,46}.
45

46
47 To induce LID, L-dopa methyl ester and benserazide were administered daily (between 9.00 and
48
49 10.00 AM) for 3 weeks at 25 and 5 mg/kg, s.c., respectively, starting from the fourth week post-
50
51 surgery ^{47,48}. This dose of L-dopa is the same as typically used in the behavioral sensitization paradigm
52
53 ⁴⁹⁻⁵⁴ and this type of sensitized context was associated with LID in PD patients ⁶. Amantadine or its
54
55 vehicle was administered at 20, 30 and 40 mg/kg, i.p., 40 min before L-dopa/benserazide
56
57 administration ^{16,55,56}.
58
59
60

Behavioral procedures

Cylinder test

Forelimb use asymmetry was assessed as previously described⁵⁷ 3 weeks after cerebral injection of 6-OHDA. Briefly, animals were placed in a glass cylinder (approximately 20 cm in diameter and 35 cm high) and video-recorded for 10 minutes to allow retrospective analysis of the behavior. A blinded observer scored the number of contacts made by individual forepaws with the cylinder wall. The percentage of left paw touches out of a total of 20 touches was determined. Limb use asymmetry was demonstrated by expressing the use of the impaired paw as a percentage of the total number of touches, with unbiased animals having a score of 50%.

Stepping test

Forelimb akinesia was assessed as previously described⁵⁸ 3 weeks after cerebral injection of 6-OHDA. Briefly, the experimenter firmly held the rat's hindquarters while it supported its weight on its contralateral forelimb. Then, the experimenter moved the rat forward along the table (0.9 m in 5 seconds) three consecutive times per session. All sessions were video-recorded and the number of adjusting steps was counted afterwards by a blinded investigator. For each session, the total score calculated was the sum of the number of adjusting steps observed for the contralateral paw in the three tests. The sessions took place between 10.30 and 11.30 AM, 3 weeks post-surgery.

L-dopa induced dyskinesia

L-dopa mediated induction of AIMs is only possible in animals with a massive nigral DA lesion^{26,59}. Thus, the 8 rats displaying a full DA lesion (i.e., over 90 contralateral turns after apomorphine challenge^{60,61}) were included in this procedure. These rats received a daily injection of L-dopa and benserazide for 3 weeks (15 testing sessions), starting the 4th week post-surgery (after the stepping test and cylinder test procedures). Each session was performed between 9.00 and 10.00 AM.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Rats were placed in a cylinder (20 cm in diameter and 35 cm high) 20 min after the L-dopa/benserazide administration and their behavior was video-recorded for 1 hour in order to allow off-line scoring of the LID by an experimenter who was unaware of the treatment condition.

Then, 4 additional testing sessions were performed in order to assess the efficacy of amantadine in reversing the LID. Animals received vehicle on day 1, amantadine 40 mg/kg on day 2, amantadine 30 mg/kg on day 3 and amantadine 20 mg/kg on day 4 (**Fig. 1**). Amantadine or its vehicle was administered 40 min before L-dopa/benserazide administration and 1 hour before the animals were placed in the cylinder and video-tracked for 1 hour. Again, the behavior of the animals was video-recorded and LID scoring was performed retrospectively by a blinded experimenter.

Consistent with previous reports that assumed different levels of severity depending on the features of the AIMS¹⁷⁻¹⁹, we hypothesized that different behaviors reflected different LID intensity levels. We attributed a global severity score to each rat, based on both the frequency and the intensity of the behaviors, in order to reflect gradual levels of AIM severity. The AIMS were classified into 5 subtypes according to their level of severity according to previously published rating scales^{14,16-22} (see **Table 1**). The purpose of our approach was to simplify the scoring method and provide a global AIM score reflecting the dyskinetic state of each animal. To this end, we developed a rating scale including both axial dystonia (neck/trunk torsion toward the side contralateral to the lesion) and forelimb dyskinesia (side to side, up to down tapping or circular movement of the right forelimb, “grabbing”) within the same assessment grid. For instance, the AIM subtypes “increased locomotion with grabbing” and “axial posture” include both changes in neck/trunk position and grabbing behavior (**Table 1**). In line with previous scales^{14,16}, contralateral circling (included under the item “increased locomotion with contralateral bias” in **Table 1**) was not considered “dyskinesia,” since contralateral turning can be induced by dopaminergic agonists with very low dyskinesiogenic potential^{16,62}. This behavior was considered “normal behavior” following L-dopa administration in unilaterally 6-OHDA lesioned rats

and, thus, corresponds to the lowest severity score. The purpose was to eventually weight the score if AIMs such as grabbing or others described in **Table 1** were observed.

Each rat was observed for 1 min every 10 min, from 20 to 80 min post L-dopa/benserazide administration. For each subtype of behavior (**Table 1**), a frequency score between 0 and 3 (with 0 = absent, 1 = occasional: less than 50% of the time, 2 = frequent: more than 50% of the time, 3 = continuous, uninterrupted) was assigned.

These frequency scores were multiplied by a severity coefficient (see **Table 1**) that was determined prior to the behavioral analysis. These severity scores were adapted from previous studies applying similar amplitude scores^{17,18,63–65} in order to maximize the sensitivity of the scale and to reflect as much contrast as possible within the different levels of AIM severity. In our study, severity coefficients were applied as follows: **1** = horizontal body position, contralateral circling on 4 paws, **2** = horizontal body position, contralateral circling on 4 paws, grabbing movement with contralateral forelimb paw, **8** = horizontal body position with pronounced deviation toward the contralateral side (>30°) with nose close to the level of the tail, possible loss of balance due to the twisted position, **12** = vertical body position, contralateral deviation of the head, neck and upper trunk 60°-90°, grabbing and contralateral rotations on hindlimbs, **20** = vertical body position with sustained and severe torsion of neck and trunk at 90°-180° causing the rat to lose balance, purposeless "choreiform" twisting movements, animal stuck in the twisted position/possible contralateral rotations on hindlimbs. The 2 bounds coefficients (1 and 20) were determined by taking into account the maximum level of dyskinetic state of the highest dyskinetic rat (i.e. rat stuck in 180° twisted position for the entire minute of observation) after 3 weeks of L-dopa treatment and the minimum level of dyskinetic state of the lowest dyskinetic rat (i.e. classical contralateral circling with no AIM during the minute of observation) on the first day of the 3-week period of L-dopa treatment. This first step aims at calibrating the system. The intermediary coefficients (2, 8 and 12) were determined arbitrarily but in order to discriminate well between the different AIM severity levels. Thus, we applied a moderate

factor of magnitude between “increased locomotion with contralateral bias” and “increased locomotion with grabbing” (1 to 2), then a more important difference between “increased locomotion with grabbing” and “dystonic circling” (2 to 8), a mild to slight difference between behaviors in which a pronounced deviation of the body is observed (8, 12 and 20 for “dystonic circling” and “axial posture and “dystonia” respectively) and a substantial factor of magnitude between “increased locomotion with contralateral bias” and “dystonia” (1 to 20).

For animals displaying different subtypes of behaviors during the same observation period, an adjustment coefficient was also applied, as described in **Table 1**. This adjustment coefficient aimed to improve the accuracy of the scoring method by taking into account slight behavioral differences. For instance, we assumed that a rat displaying only “dystonia” (see **Table 1**) during the entire minute of observation time was more severely affected than a rat that displayed “dystonia” most of the time but that also displayed “axial posture” and/or other less severe symptoms.

A global severity score taking into account both the frequency and the intensity of the AIMS was obtained for each rat (the detailed calculations and formulae used are available in **Supplementary Material 1**). The maximum score that could be accumulated per testing session was 420 (maximum score per observation point: 60; number of observation points per session: 7).

Histological analysis

After the last day of LID scoring, the animals were decapitated under pentobarbital anesthesia (200 mg/kg) and the brain was quickly removed at 4°C on fresh ice. The brains were then post-fixed in ice-cold 4% paraformaldehyde over 24h and processed for paraffin embedding before being sectioned (7-µm slices) in the coronal plane. The slides were stained using antibodies against tyrosine hydroxylase (TH) (rabbit 1/5000: Millipore AB152) and the DAKO LSAB and HRP system (DAKO Real, DAKO France). Immunostaining was processed on a Discovery XT® Platform (Roche Ventana, Tucson,

AZ, USA). All slides were scanned at 20x magnification for whole slide imaging using a NanoZoomer 2.0RS scanner (Hamamatsu, Japan) and analyzed using NIS-AR software (Nikon, Japan).

TH+ cells in the substantia nigra pars compacta (SNc) were counted at 3 different rostrocaudal levels (AP: -4.80 mm, -5.30 mm and -6.04 mm) according to the Paxinos rat brain atlas⁴². At each level, the number of TH+ cells contained in the SNc was counted in three adjacent sections by a blinded experimenter. For each section, the boundaries were chosen by examining the shape of the cells and referring to the Paxinos rat brain atlas⁴². At 100x magnification, only the cells with a pyramidal shape were counted. For each animal, the total number of TH+ cells was estimated using the Königsmark formula: $N_t = N_s \times (S_t/S_s)$ where N_t =total number of cells; N_s =number of cells counted; S_t =total number of sections counted; S_s =total number of sections through the SNc^{58,66}. The percentage of DA cell loss was estimated using the following formula: $100 - (TH+ \text{ cell number in lesioned SNc} / TH+ \text{ cells number in sham SNc} \times 100)$.

Data and statistical analysis

Data were analyzed by Student's t-test, one-way ANOVA, repeated measures ANOVA or two-way repeated measures ANOVA depending on the experimental design. Where appropriate, post-hoc analyses were carried out with Tukey's or Dunnett's tests. All reported p-values are two-sided. The normality of the variables was assessed by Shapiro-Wilk test and the homoscedasticity was assessed by Levene's test. When data were not normally distributed, the data were square-transformed. Results were expressed as means $\pm$ SEM.

Supporting Information

Excel sheet containing all formulae for LID score calculation.

Author Contributions

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

SL designed the study, analyzed the data and wrote the manuscript. ASD, WD, MA, PM and MS performed the experiments and CDLR reviewed the manuscript.

Funding Sources

This research was funded by Biotrial Pharmacology.

Conflict of Interest

SL, ASD, WD, MA, PM and CDLR are employees of Biotrial Pharmacology.

Acknowledgements

The authors thank Cliona McSweeney, Fiona McAlpine and Stéphanie Le Goaller for providing very useful comments and for reviewing the manuscript.

References

1. Cenci, M. A., Ohlin, K. E. & Rylander, D. Plastic effects of L-DOPA treatment in the basal ganglia and their relevance to the development of dyskinesia. *Park. Relat Disord* **15 Suppl 3**, S59-63 (2009).
2. LeWitt, P. A. Levodopa therapy for Parkinson's disease: Pharmacokinetics and pharmacodynamics. *Mov Disord* **30**, 64–72 (2015).
3. Ahlskog, J. E. & Muenter, M. D. Frequency of levodopa-related dyskinesias and motor fluctuations as estimated from the cumulative literature. *Mov. Disord. Off. J. Mov. Disord. Soc.* **16**, 448–458 (2001).
4. Grandas, F., Galiano, M. L. & Tabernero, C. Risk factors for levodopa-induced dyskinesias in Parkinson's disease. *J. Neurol.* **246**, 1127–1133 (1999).
5. Iderberg, H., Francardo, V. & Pioli, E. Y. Animal models of L-DOPA-induced dyskinesia: an update on the current options. *Neuroscience* **211**, 13–27 (2012).
6. Bezard, E., Brotchie, J. M. & Gross, C. E. Pathophysiology of levodopa-induced dyskinesia: potential for new therapies. *Nat. Rev. Neurosci.* **2**, 577–588 (2001).
7. Buck, K. & Ferger, B. L-DOPA-induced dyskinesia in Parkinson's disease: a drug discovery perspective. *Drug Discov. Today* **15**, 867–875 (2010).
8. Meissner, W. G. *et al.* Priorities in Parkinson's disease research. *Nat. Rev. Drug Discov.* **10**, 377–393 (2011).
9. Schapira, A. H. V., Olanow, C. W., Greenamyre, J. T. & Bezard, E. Slowing of neurodegeneration in Parkinson's disease and Huntington's disease: future therapeutic perspectives. *Lancet* **384**, 545–555 (2014).
10. Carta, A. R., Pinna, A. & Morelli, M. How reliable is the behavioural evaluation of dyskinesia in animal models of Parkinson's disease? *Behav. Pharmacol.* **17**, 393–402 (2006).
11. Morin, N., Jourdain, V. A. & Di Paolo, T. Modeling dyskinesia in animal models of Parkinson disease. *Exp. Neurol.* **256**, 105–116 (2014).

12. Kola, I. The state of innovation in drug development. *Clin Pharmacol Ther* **83**, 227–30 (2008).
13. Kola, I. & Landis, J. Can the pharmaceutical industry reduce attrition rates? *Nat Rev Drug Discov* **3**, 711–5 (2004).
14. Cenci, M. A., Lee, C. S. & Björklund, A. L-DOPA-induced dyskinesia in the rat is associated with striatal overexpression of prodynorphin- and glutamic acid decarboxylase mRNA. *Eur. J. Neurosci.* **10**, 2694–2706 (1998).
15. Duty, S. & Jenner, P. Animal models of Parkinson's disease: a source of novel treatments and clues to the cause of the disease: Animal models of Parkinson's disease. *Br. J. Pharmacol.* **164**, 1357–1391 (2011).
16. Lundblad, M. *et al.* Pharmacological validation of behavioural measures of akinesia and dyskinesia in a rat model of Parkinson's disease. *Eur. J. Neurosci.* **15**, 120–132 (2002).
17. Cenci, M. A. & Lundblad, M. Ratings of L-DOPA-Induced Dyskinesia in the Unilateral 6-OHDA Lesion Model of Parkinson's Disease in Rats and Mice. in *Current Protocols in Neuroscience* (eds. Crawley, J. N. *et al.*) (John Wiley & Sons, Inc., 2007).
18. Winkler, C., Kirik, D., Björklund, A. & Cenci, M. A. L-DOPA-Induced Dyskinesia in the Intrastriatal 6-Hydroxydopamine Model of Parkinson's Disease: Relation to Motor and Cellular Parameters of Nigrostriatal Function. *Neurobiol. Dis.* **10**, 165–186 (2002).
19. Steece-Collier, K. *et al.* Embryonic mesencephalic grafts increase levodopa-induced forelimb hyperkinesia in parkinsonian rats. *Mov. Disord.* **18**, 1442–1454 (2003).
20. Breger, L. S., Dunnett, S. B. & Lane, E. L. Comparison of rating scales used to evaluate L-DOPA-induced dyskinesia in the 6-OHDA lesioned rat. *Neurobiol. Dis.* **50**, 142–150 (2013).
21. Dekundy, A., Lundblad, M., Danysz, W. & Cenci, M. A. Modulation of L-DOPA-induced abnormal involuntary movements by clinically tested compounds: Further validation of the rat dyskinesia model. *Behav. Brain Res.* **179**, 76–89 (2007).

22. Michel, A., Christophe, B. & Scheller, D. Comparison of the activity of the three NR2B antagonists, Ro 25-6981, Co 101244 and 20J in experimental models of Parkinson's disease. in 742.8/U17 (Society for Neuroscience, 2008).
23. Johnson, G. S. Commercial viability of CNS drugs: balancing the risk/reward profile. *Neurobiol. Dis.* **61**, 21–24 (2014).
24. Pangalos, M. N., Schechter, L. E. & Hurko, O. Drug development for CNS disorders: strategies for balancing risk and reducing attrition. *Nat. Rev. Drug Discov.* **6**, 521–532 (2007).
25. Lane, E. & Dunnett, S. Animal models of Parkinson's disease and L-dopa induced dyskinesia: how close are we to the clinic? *Psychopharmacology (Berl.)* **199**, 303–312 (2008).
26. Nadjar, A., Gerfen, C. R. & Bezard, E. Priming for l-dopa-induced dyskinesia in Parkinson's disease: a feature inherent to the treatment or the disease? *Prog. Neurobiol.* **87**, 1–9 (2009).
27. Colosimo, C. *et al.* Task force report on scales to assess dyskinesia in Parkinson's disease: critique and recommendations. *Mov. Disord. Off. J. Mov. Disord. Soc.* **25**, 1131–1142 (2010).
28. Müller, T. New small molecules for the treatment of Parkinson's disease. *Expert Opin. Investig. Drugs* **19**, 1077–1086 (2010).
29. Bastide, M. F. *et al.* Immediate-early gene expression in structures outside the basal ganglia is associated to l-DOPA-induced dyskinesia. *Neurobiol. Dis.* **62**, 179–192 (2014).
30. Charbonnier-Beaupel, F. *et al.* Gene expression analyses identify Narp contribution in the development of L-DOPA-induced dyskinesia. *J. Neurosci. Off. J. Soc. Neurosci.* **35**, 96–111 (2015).
31. Grünblatt, E., Schmidt, W. J., Scheller, D. K. A., Riederer, P. & Gerlach, M. Transcriptional alterations under continuous or pulsatile dopaminergic treatment in dyskinetic rats. *J. Neural Transm. Vienna Austria 1996* **118**, 1717–1725 (2011).
32. Andersen, A. D. *et al.* Cerebrospinal fluid levels of catecholamines and its metabolites in Parkinson's disease: effect of l-DOPA treatment and changes in levodopa-induced dyskinesia. *J. Neurochem.* (2017). doi:10.1111/jnc.13997

33. López-Azcárate, J. *et al.* Coupling between beta and high-frequency activity in the human subthalamic nucleus may be a pathophysiological mechanism in Parkinson's disease. *J. Neurosci. Off. J. Soc. Neurosci.* **30**, 6667–6677 (2010).
34. Litvak, V. *et al.* Movement-related changes in local and long-range synchronization in Parkinson's disease revealed by simultaneous magnetoencephalography and intracranial recordings. *J. Neurosci. Off. J. Soc. Neurosci.* **32**, 10541–10553 (2012).
35. Jenkinson, N., Kühn, A. A. & Brown, P. γ oscillations in the human basal ganglia. *Exp. Neurol.* **245**, 72–76 (2013).
36. Dupre, K. B. *et al.* Effects of L-dopa priming on cortical high beta and high gamma oscillatory activity in a rodent model of Parkinson's disease. *Neurobiol. Dis.* **86**, 1–15 (2016).
37. Aristieta, A., Ruiz-Ortega, J. A., Miguelez, C., Morera-Herreras, T. & Ugedo, L. Chronic L-DOPA administration increases the firing rate but does not reverse enhanced slow frequency oscillatory activity and synchronization in substantia nigra pars reticulata neurons from 6-hydroxydopamine-lesioned rats. *Neurobiol. Dis.* **89**, 88–100 (2016).
38. Miller, G. Is pharma running out of brainy ideas? *Science* **329**, 502–504 (2010).
39. Research and development: Outsourcing trends. *Nature* **512**, 106–106 (2014).
40. Wegener, G. & Rujescu, D. The current development of CNS drug research. *Int. J. Neuropsychopharmacol.* **16**, 1687–1693 (2013).
41. Yokley, B. H., Hartman, M. & Slusher, B. S. Role of Academic Drug Discovery in the Quest for New CNS Therapeutics. *ACS Chem. Neurosci.* **8**, 429–431 (2017).
42. Paxinos, G. & Watson, C. *The Rat Brain in Stereotaxic Coordinates*. (Academic Press, 1998).
43. Michel, A. *et al.* Behavioural Assessment of the A2a/NR2B Combination in the Unilateral 6-OHDA-Lesioned Rat Model: A New Method to Examine the Therapeutic Potential of Non-Dopaminergic Drugs. *PLOS ONE* **10**, e0135949 (2015).

44. Metz, G. A., Tse, A., Ballermann, M., Smith, L. K. & Fouad, K. The unilateral 6-OHDA rat model of Parkinson's disease revisited: an electromyographic and behavioural analysis. *Eur. J. Neurosci.* **22**, 735–744 (2005).
45. Kelsey, J. E. *et al.* NMDA receptor antagonists ameliorate the stepping deficits produced by unilateral medial forebrain bundle injections of 6-OHDA in rats. *Psychopharmacology (Berl.)* **175**, 179–188 (2004).
46. Tronci, E. *et al.* 5-Hydroxy-tryptophan for the treatment of L-DOPA-induced dyskinesia in the rat Parkinson's disease model. *Neurobiol. Dis.* **60**, 108–114 (2013).
47. Xie, C. *et al.* Inhibition of Glycogen Synthase Kinase-3 β (GSK-3 β) as potent therapeutic strategy to ameliorates L-dopa-induced dyskinesia in 6-OHDA parkinsonian rats. *Sci. Rep.* **6**, 23527 (2016).
48. Zhang, S., Xie, C., Wang, Q. & Liu, Z. Interactions of CaMKII with dopamine D2 receptors: roles in levodopa-induced dyskinesia in 6-hydroxydopamine lesioned Parkinson's rats. *Sci. Rep.* **4**, 6811 (2014).
49. Bordet, R. *et al.* Induction of dopamine D3 receptor expression as a mechanism of behavioral sensitization to levodopa. *Proc. Natl. Acad. Sci. U. S. A.* **94**, 3363–3367 (1997).
50. Bordet, R., Ridray, S., Schwartz, J. C. & Sokoloff, P. Involvement of the direct striatonigral pathway in levodopa-induced sensitization in 6-hydroxydopamine-lesioned rats. *Eur. J. Neurosci.* **12**, 2117–2123 (2000).
51. Brown, A. M., Deutch, A. Y. & Colbran, R. J. Dopamine depletion alters phosphorylation of striatal proteins in a model of Parkinsonism. *Eur. J. Neurosci.* **22**, 247–256 (2005).
52. Bychkov, E., Ahmed, M. R., Dalby, K. N. & Gurevich, E. V. Dopamine depletion and subsequent treatment with L-DOPA, but not the long-lived dopamine agonist pergolide, enhances activity of the Akt pathway in the rat striatum. *J. Neurochem.* **102**, 699–711 (2007).
53. van Kampen, J. M. & Stoessl, A. J. Effects of oligonucleotide antisense to dopamine D3 receptor mRNA in a rodent model of behavioural sensitization to levodopa. *Neuroscience* **116**, 307–314 (2003).

- 1
2
3 54. Sgambato-Faure, V. *et al.* Coordinated and spatial upregulation of arc in striatonigral neurons
4
5 correlates with L-dopa-induced behavioral sensitization in dyskinetic rats. *J. Neuropathol. Exp.*
6
7 *Neurol.* **64**, 936–947 (2005).
8
- 9 55. Bido, S., Marti, M. & Morari, M. Amantadine attenuates levodopa-induced dyskinesia in mice
10
11 and rats preventing the accompanying rise in nigral GABA levels: Nigral GABA in dyskinetic mice
12
13 and rats. *J. Neurochem.* **118**, 1043–1055 (2011).
14
- 15 56. Bortolanza, M., Bariotto-Dos-Santos, K. D., Dos-Santos-Pereira, M., da-Silva, C. A. & Del-Bel, E.
16
17 Antidyskinetic Effect of 7-Nitroindazole and Sodium Nitroprusside Associated with Amantadine
18
19 in a Rat Model of Parkinson's Disease. *Neurotox. Res.* **30**, 88–100 (2016).
20
- 21 57. Decressac, M., Mattsson, B. & Björklund, A. Comparison of the behavioural and histological
22
23 characteristics of the 6-OHDA and α -synuclein rat models of Parkinson's disease. *Exp. Neurol.*
24
25 **235**, 306–315 (2012).
26
- 27 58. Paillé, V., Henry, V., Lescaudron, L., Brachet, P. & Damier, P. Rat model of Parkinson's disease
28
29 with bilateral motor abnormalities, reversible with levodopa, and dyskinesias. *Mov. Disord.* **22**,
30
31 533–539 (2007).
32
33
- 34 59. Blandini, F., Armentero, M.-T. & Martignoni, E. The 6-hydroxydopamine model: News from the
35
36 past. *Parkinsonism Relat. Disord.* **14**, S124–S129 (2008).
37
- 38 60. Papa, S. M., Engber, T. M., Kask, A. M. & Chase, T. N. Motor fluctuations in levodopa treated
39
40 parkinsonian rats: relation to lesion extent and treatment duration. *Brain Res.* **662**, 69–74 (1994).
41
42
- 43 61. Kaakkola, S. & Teräväinen, H. Animal models of parkinsonism. *Pharmacol. Toxicol.* **67**, 95–100
44
45 (1990).
46
- 47 62. Carta, A. R., Frau, L., Pontis, S., Pinna, A. & Morelli, M. Direct and indirect striatal efferent
48
49 pathways are differentially influenced by low and high dyskinetic drugs: behavioural and
50
51 biochemical evidence. *Parkinsonism Relat. Disord.* **14 Suppl 2**, S165-168 (2008).
52
53
54
55
56
57
58
59
60

- 1
2
3 63. Lane, E. L., Soulet, D., Vercammen, L., Cenci, M. A. & Brundin, P. Neuroinflammation in the
4
5 generation of post-transplantation dyskinesia in Parkinson's disease. *Neurobiol. Dis.* **32**, 220–228
6
7 (2008).
8
9 64. Mela, F. *et al.* Antagonism of metabotropic glutamate receptor type 5 attenuates L-DOPA-
10
11 induced dyskinesia and its molecular and neurochemical correlates in a rat model of Parkinson's
12
13 disease. *J. Neurochem.* **101**, 483–497 (2007).
14
15 65. Monville, C., Torres, E. M., Pekarik, V., Lane, E. L. & Dunnett, S. B. Genetic, temporal and diurnal
16
17 influences on L-dopa-induced dyskinesia in the 6-OHDA model. *Brain Res. Bull.* **78**, 248–253
18
19 (2009).
20
21 66. German, D. C. & Manaye, K. F. Midbrain dopaminergic neurons (nuclei A8, A9, and A10): three-
22
23 dimensional reconstruction in the rat. *J Comp Neurol* **331**, 297–309 (1993).
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table Legends

Table 1. Description and severity classification of the AIMs observed. Severity and adjustment coefficients were applied according to the type of AIMs observed.

AIM subtype	Increased locomotion with contralateral bias	Increased locomotion with grabbing	Dystonic circling	Axial posture	Dystonia
Description of the behavior	horizontal body position contralateral circling on 4 paws	horizontal body position contralateral circling on 3 paws grabbing movement with contralateral forelimb paw	horizontal body position with pronounced deviation toward the contralateral side nose close to the level of the tail possible loss of balance due to the twisted position	vertical body position contralateral deviation of the head, neck and trunk grabbing and contralateral rotations on hindlimbs	vertical body position with sustained and severe 180° trunk torsion purposeless "choreiform" twisting movements, loss of balance, falls animal stuck in the twisted position/possible contralateral rotations on hindlimbs
Severity coefficient	1	2	8	12	20
Adjustment coefficient	-0,6	-0,4	-0,2	-0,1	none

Figure Legends

Fig. 1. Study design. General experimental procedure and timeline.

Fig. 2. Assessment of nigral loss and forelimb use asymmetry and akinesia. (a) Immunolabeling of tyrosine hydroxylase (TH). The bar corresponds to 1 mm. VTA = ventral tegmental area (b) Number of TH+ cells counted in the SNc (n=18). An unpaired Student's t-test revealed severe DA cell loss in the SNc on the lesioned side compared to the unlesioned side ($t_{34}=11.09$; $***p<0.0001$). All rats displayed a lesion with over 80% cell loss compared to the unlesioned side, except for 2 rats with partial lesions with 31% and 44% cell loss, respectively (black dots). Center lines and error bars throughout the figure represent means $\pm$ SEM. (c) Contralateral paw use in the cylinder test (n=8-10). A one-way ANOVA revealed a significant treatment effect, with $F_{2,23}=15.6$; $p<0.0001$. The percentage of contralateral paw use was significantly decreased in vehicle-treated lesioned rats compared to sham rats ($***p=0.0005$ with Tukey's post-hoc test) with L-dopa/benserazide (8 and 6 mg/kg, ip respectively) reversing this impairment (no significant difference compared to sham rats and $***p<0.0001$ when compared to lesioned vehicle-treated rats with Tukey's post-hoc test). (d) Number of adjusting steps performed with the contralateral paw (n=8-10). A one-way ANOVA revealed a significant treatment effect, with $F_{2,23}=42.21$; $p<0.0001$. The number of adjusting steps was significantly decreased in vehicle-treated lesioned rats compared to sham rats ($***p<0.0001$ with Tukey's post-hoc test) with L-dopa/benserazide (8 and 6 mg/kg, ip respectively) alleviating this impairment ($**p=0.0057$ when compared to sham rats and $***p<0.0001$ when compared to vehicle-treated lesioned rats with Tukey's post-hoc test). (e) Apomorphine-induced contralateral rotations (n=8-18). An unpaired Student's t-test revealed that lesioned rats produced significantly more contralateral rotations compared to sham ($t_{24}=2.183$; $p=0.0391$), with 8 rats displaying over 90 turns/45 min (dotted line). Ten other lesioned rats were below this threshold (the 2 black dots represent the partially lesioned rats). Center lines and error bars throughout the figure represent means $\pm$ SEM.

Fig. 3. L-dopa induced dyskinesia. (a) Time course of the development of LID over 3 weeks of daily L-dopa/benserazide treatment (n=8). The ranked data were square-transformed to allow normal distribution and the use of parametric statistics. A repeated measures one-way ANOVA revealed a significant time effect, with $F_{1.549,10.84}=16.6$; $p=0.0009$. The data reflect a time-dependent increase in the LID score over time with a significant increase in week 1, week 2 and week 3 compared to week 0

(*p=0.026; *p=0.0387 and *p=0.0362, respectively, with Dunnett's post-hoc test). **(b)**. Contribution of the different AIM components to the global LID score over the 3 weeks of L-dopa treatment (n=8). Adjustment coefficients were not applied (as their purpose is to weight the global LID score) and the ranked data were square-transformed to allow normal distribution and the use of parametric statistics (but untransformed data are displayed to improve clarity). For each time-point, a one way ANOVA followed by a Tukey's post-hoc tests revealed a significant increase of "increased locomotion with contralateral bias" and "dystonia" at week 0 and week 3 respectively (**p<0.001 and *p<0.05 respectively) compared to other behavioral subtypes. A one-way ANOVA revealed a significant time effect for "increased locomotion with grabbing" ($F_{3,28}=9.056$; $p=0.0002$), "axial posture" ($F_{3,28}=7.594$; $p=0.0007$) and "dystonia" ($F_{3,28}=4.287$; $p=0.0131$). A Dunnett's post-hoc analysis showed a significant increase of "increased locomotion with grabbing" on week 1 and week 2 compared to week 0 (***p=0.0008 and **p=0.0038 respectively), a significant increase of "axial posture" and "dystonia" on week 3 compared to week 0 (***p=0.0009 and *p=0.0151 respectively). Turning = increased locomotion with contralateral bias, Turning+Grab. = increased locomotion with grabbing, Dyst. Circling = dystonic circling. **(c)** Amantadine-mediated alleviation of the LID score (n=8). The ranked data were square-transformed to allow normal distribution and the use of parametric statistics. A repeated measures one-way ANOVA revealed a significant treatment effect, with $F_{1,167,8,168}=9.447$; $p=0.0129$. The data reflect a dose-dependent decrease in the LID score after treatment with amantadine, with the maximum effect at 40 mg/kg (**p=0.0034 compared to vehicle with Dunnett's post-hoc test). **(d)** Time course of the effect of amantadine (n=8). The ranked data were square-transformed to allow normal distribution and the use of parametric statistics. A repeated measures two-way ANOVA revealed no significant treatment effect ($F_{3,28}=2.609$; $p=0.0712$), but a significant time effect ($F_{6,168}=8.9$; $p<0.0001$) and a significant time x treatment interaction ($F_{18,168}=1.756$; $p=0.0345$). The Dunnett's post-hoc analysis showed a significant effect of amantadine 40 mg/kg compared to vehicle at 30 min (**p=0.007), 40 min (**p=0.0019), 50 min (**p=0.002) and 70 min (*p=0.0155) post-L-dopa administration. The 30 mg/kg dose also significantly decreased the LID score compared to vehicle at 50 min post-L-dopa administration (*p=0.0443 with Dunnett's post-hoc test). Ama20, Ama30 and Ama40 = amantadine 20 mg/kg, 30 mg/kg and 40 mg/kg respectively. **(e)** Individual effect of amantadine 40 mg/kg on LID score (n=8). The ranked data were square-transformed to allow normal distribution and the use of parametric statistics. A paired Student's t test revealed a significant decrease in the LID score compared to vehicle-treated rats ($t_7=3.233$; $p=0.0144$). **(f)** Effect of amantadine on the different behavioral subtypes. Adjustment coefficients were not applied (as their purpose is to weight the global LID score) and the ranked data were square-transformed to allow normal distribution and the use of parametric statistics (but untransformed data are displayed to improve clarity). For each group of treatment, a one way

1
2
3 ANOVA followed by a Tukey's post-hoc tests revealed a significant increase of "dystonia" and "axial
4 posture" scores compared to other behavioral subtypes (* $p < 0.05$; ** $p < 0.01$ and *** $p < 0.001$
5 respectively). A one-way ANOVA revealed a significant treatment effect for the dystonia component
6 ($F_{3,28} = 3.066$; $p = 0.0442$). A Dunnett's post-hoc analysis showed a significant effect of amantadine 30
7 mg/kg and 40 mg/kg compared to vehicle ($^{\#}p = 0.0379$ and $^{\#}p = 0.035$ respectively) for this particular
8 behavioral subtype.
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

FIGURE 1

FIGURE 2

FIGURE 3

For Table of Contents Use Only

Validation of a new scoring scale for behavioral assessment of L-dopa-induced dyskinesia in the rat: a new tool for early decision-making in drug development

Simon Loiodice, Anne-Sophie Denibaud, Wendy Deffains, Magali Alix, Pierre Montagne, Marine Seffals, Christophe Drieu La Rochelle