Survey on the p -adic Hayman conjecture

by Alain Escassut and Jacqueline Ojeda

Introduction

We denote by $\mathbb{K}$ an algebraically closed field of characteristic 0, complete with respect to an ultrametric absolute value $|\cdot|$ and by $\mathbb{K}(x)$ the field of rational functions with coefficients in $\mathbb{K}$. Given $a \in \mathbb{K}$ and $R \in \mathbb{R}_+^*$, we denote by $d(a, R)$ the disk $\{x \in \mathbb{K} \mid |x - a| \leq R\}$, by $d(a, R^-)$ the disk $\{x \in \mathbb{K} \mid |x - a| < R\}$ and by $C(a, r)$ the circle $\{x \in \mathbb{K} \mid |x - a| = r\}$. Next, we denote by $\mathcal{A}(\mathbb{K})$ the $\mathbb{K}$ -algebra of analytic functions in $\mathbb{K}$ (i.e. the set of power series with an infinite radius of convergence) and by $\mathcal{M}(\mathbb{K})$ the field of meromorphic functions in $\mathbb{K}$ (i.e. the field of fractions of $\mathcal{A}(\mathbb{K})$). Similarly, we denote by $\mathcal{A}(d(a, R^-))$ the $\mathbb{K}$ -algebra of analytic functions in $d(a, R^-)$ (i.e. the set of power series with a radius of convergence $\geq R$) [8], [11], [12], [13] and by $\mathcal{M}(d(a, R^-))$ its field of fractions and we denote by $\mathcal{A}_b(d(a, R^-))$ the $\mathbb{K}$ -algebra of bounded analytic functions in $d(a, R^-)$ and by $\mathcal{M}_b(d(a, R^-))$ its field of fractions and we set $\mathcal{A}_u(d(a, R^-)) = \mathcal{A}(d(a, R^-)) \setminus \mathcal{A}_b(d(a, R^-))$ and $\mathcal{M}_u(d(a, R^-)) = \mathcal{M}(d(a, R^-)) \setminus \mathcal{M}_b(d(a, R^-))$.

Given $f \in \mathcal{M}(\mathbb{K})$ (resp. $f \in \mathcal{M}_u(d(a, R^-))$), a value $b \in \mathbb{K}$ is called *an exceptional value for f* if $f - b$ has no zero in $\mathbb{K}$ (resp. in $d(a, R^-)$) and it is called *a quasi-exceptional value for f* if $f - b$ has finitely many zeros in $\mathbb{K}$ (resp. in $d(a, R^-)$).

In the complex field, in the fifties, Walter Hayman asked the question whether, given a meromorphic function g in the whole plane $\mathbb{C}$ and an integer $n \in \mathbb{N}$, the function $g'g^n$ might admit an exceptional value $b \neq 0$ or a quasi-exceptional value $b \neq 0$ [10]. W. Hayman showed that $g'g^n$ has no quasi-exceptional value, whenever $n \geq 3$. Next, the problem was solved for $n = 2$ by E. Mues in 1979 [15] and next, for $n = 1$, in 1995 by W. Bergweiler and A. Eremenko [1] and separately by H. Chen and M. Fang.

The same problem occurs on the field $\mathbb{K}$, both in $\mathcal{M}(\mathbb{K})$ and in a field $\mathcal{M}(d(a, R^-))$, $a \in \mathbb{K}$, $R > 0$. Several basic results will be necessary to examine this.

In ultrametric analysis as in complex analysis, we have this immediate correspondance:

Lemma 1: *Let $g \in \mathcal{M}(\mathbb{K})$ (resp. let $g \in \mathcal{M}(d(a, R^-))$, $a \in \mathbb{K}$, $R > 0$), let $f = \frac{1}{g}$ and let $n \in \mathbb{N}^*$. Then $g'g^n$ admits a quasi-exceptional value $b \in \mathbb{K}^*$ if and only if $f' + bf^{n+2}$ has finitely many zeros that are not zeros of f .*

Remark: We can also consider the same problem when $n = -1$ i.e. the question whether $f' + bf$ has infinitely many zeros. In $\mathbb{C}$ the well known counter-example furnished by the function $\exp(x)$ shows that $f' + f$ may have no zero. When $n = 0$, in $\mathbb{C}$ the well known counter-example furnished by the function $\tan(-x)$ shows that $f' + f^2$ may have no zero. On the field $\mathbb{K}$, we will examine the cases $n = -1$ and $n = 0$.

Henceforth, for $n \geq 3$, we will examine that problem by considering the set of zeros of $f' + bf^{n+2}$, with $b \neq 0$. In the field $\mathbb{K}$, two theorems are specific to p-adic analysis. Both are based on the following lemma. We set $m = n + 2$.

Notation: Let $f \in \mathcal{M}(d(0, R^-))$. For every $r \in]0, R[$, $|f(x)|$ has a limit when $|x|$ tends to r while staying different from r and that limit is denoted by $|f|(r)$ [8].

Let $\mu = \log r$. We set $\Psi(f, \mu) = \log(|f|(r))$. We denote by $\nu^+(f, \mu)$ the difference between the number of zeros and the number of poles of f in $d(0, r)$ and we denote by $\nu^-(f, \mu)$ the difference between the number of zeros and the number of poles of f in $d(0, r^-)$.

The following Lemma 1 and 2 are classical [17]:

Lemma 2: Let $f \in \mathcal{M}(\mathbb{K})$. Then $\nu^+(f, \mu)$ is the right side derivative of $\Psi(f, \mu)$ and $\nu^-(f, \mu)$ is the left side derivative of $\Psi(f, \mu)$.

Lemma 3: Let $f \in \mathcal{M}(\mathbb{K})$, (resp. let $f \in \mathcal{M}(d(a, R^-))$, $a \in \mathbb{K}$, $R > 0$), suppose that f admits infinitely many zeros and suppose that there exists a sequence of intervals $[r'_n, r''_n]$ such that $\lim_{n \rightarrow +\infty} r'_n = +\infty$ (resp. $\lim_{n \rightarrow +\infty} r'_n = \lim_{n \rightarrow +\infty} r''_n = R$) and such that $|(f' + f^m)|(r) = |f^m|(r) \forall r \in \bigcup_{n \in \mathbb{N}} [r'_n, r''_n]$. Let $m \in \mathbb{N}^*$ be $\neq 2$. Then $f' + f^m$ has infinitely many zeros that are not zeros of f .

Proof: Let $J = \bigcup_{n \in \mathbb{N}} [r'_n, r''_n]$. When r is big enough, we have $|f^m|(r) > |f'|(r)$ therefore

$$(1) \quad \nu^+(f' + f^m, \log r) = \nu^+(f^m, \log r), \quad \nu^-(f' + f^m, \log r) = \nu^-(f^m, \log r) \quad \forall r \in J.$$

Consequently, in each disk $d(0, r)$ with $r \in J$, f and $f' + f^m$ have the same difference between the number of zeros and poles. Now, if $m \geq 3$ the poles of $f' + f^m$ and f^m are the same taking multiplicity into account. And when $m = 1$, each pole of f is a pole of $f' + f$ with a greater order. Consequently, for each $r \in J$, the number of zeros of $f' + f^m$ in $d(0, r)$ is superior or equal to this of f^m .

Now, for each $n \in \mathbb{N}$, let s_n be the number of distinct zeros of f in $d(0, r''_n)$. Since f has infinitely many zeros, the sequence s_n is increasing and tends to $+\infty$. On the other hand, for each zero α of order u of f , either α is not a zero of $f' + f^m$ (when $u = 1$), or it is a zero of order $u - 1$. Consequently, the number of zeros of $f' + f^m$ in $d(0, r''_n)$ which are not zeros of f is at least s_n . Thus we have proved that $f' + f^m$ has infinitely many zeros that are not zeros of f .

Remark: Relation (1) above does not hold when $m = 2$ because poles of f^2 and f' may have the same order and therefore may kill each other.

In most of results, we will use the ultrametric Nevanlinna theory [33], [35]. The Nevanlinna Theory was made by Rolf Nevanlinna on complex functions [16], [10]. It

consists of defining counting functions of zeros and poles of a meromorphic function f and giving an upper bound for multiple zeros and poles of various functions $f - b$, $b \in \mathbb{C}$.

A similar theory for functions in a p-adic field was constructed by A. Boutabaa [6].

Notations: Given three functions ϕ , ψ , ζ defined in an interval $J =]a, +\infty[$ (resp. $J =]a, R[$), with values in $[0, +\infty[$, we shall write $\phi(r) \leq \psi(r) + O(\zeta(r))$ if there exists a constant $b \in \mathbb{R}$ such that $\phi(r) \leq \psi(r) + b\zeta(r)$. We shall write $\phi(r) = \psi(r) + O(\zeta(r))$ if $|\psi(r) - \phi(r)|$ is bounded by a function of the form $b\zeta(r)$.

Similarly, we shall write $\phi(r) \leq \psi(r) + o(\zeta(r))$ if there exists a function h from $J =]a, +\infty[$ (resp. from $J =]a, R[$) to $\mathbb{R}$ such that $\lim_{r \rightarrow +\infty} \frac{h(r)}{\zeta(r)} = 0$ (resp. $\lim_{r \rightarrow R} \frac{h(r)}{\zeta(r)} = 0$) and such that $\phi(r) \leq \psi(r) + h(r)$. And we shall write $\phi(r) = \psi(r) + o(\zeta(r))$ if there exists a function h from $J =]a, +\infty[$ (resp. from $J =]a, R[$) to $\mathbb{R}$ such that $\lim_{r \rightarrow +\infty} \frac{h(r)}{\zeta(r)} = 0$ (resp. $\lim_{r \rightarrow R} \frac{h(r)}{\zeta(r)} = 0$) and such that $\phi(r) = \psi(r) + h(r)$.

The p-adic Nevanlinna Theory was first stated and correctly proved by A. Boutabaa in $\mathcal{M}(\mathbb{K})$ [6]. In [7] the theory was extended to functions in $\mathcal{M}(d(0, R^-))$ by taking into account Lazard's problem [14].

Throughout the next paragraphs, we will denote by I the interval $[t, +\infty[$ and by J an interval of the form $[t, R[$ with $t > 0$.

We have to introduce the counting function of zeros and poles of f , counting or not multiplicity. Here we will choose a presentation that avoids assuming that all functions we consider admit no zero and no pole at the origin.

Definitions: We denote by $Z(r, f)$ the counting function of zeros of f in $d(0, r)$ in the following way:

Let $\sigma(r)$ be the number of distinct zeros of f in $d(0, r)$ and let (a_n) , $1 \leq n \leq \sigma(r)$ be the finite sequence of zeros of f in $d(0, r)$, of respective order s_n .

We set $Z(r, f) = \max(\omega_0(f), 0) \log r + \sum_{n=1}^{\sigma(r)} s_n (\log r - \log |a_n|)$ and so, $Z(r, f)$ is called *the counting function of zeros of f in $d(0, r)$, counting multiplicity*.

In order to define the counting function of zeros of f without multiplicity, we put $\overline{\omega}_0(f) = 0$ if $\omega_0(f) \leq 0$ and $\overline{\omega}_0(f) = 1$ if $\omega_0(f) \geq 1$.

Now, we denote by $\overline{Z}(r, f)$ the counting function of zeros of f without multiplicity:

$\overline{Z}(r, f) = \overline{\omega}_0(f) \log r + \sum_{n=1}^{\sigma(r)} (\log r - \log |a_n|)$ and so, $\overline{Z}(r, f)$ is called *the counting function of zeros of f in $d(0, r)$ ignoring multiplicity*.

In the same way, we denote by $\tau(r)$ the number of distinct poles of f in $d(0, r)$ and then, considering the finite sequence (b_n) , $1 \leq n \leq \tau(r)$ of poles of f in $d(0, r)$, with respective multiplicity order t_n , we put

$$N(r, f) = \max(-\omega_0(f), 0) \log r + \sum_{n=1}^{\tau(r)} t_n (\log r - \log |b_n|)$$
 and then $N(r, f)$ is called *the counting function of the poles of f , counting multiplicity*

Next, in order to define the counting function of poles of f without multiplicity, we put $\overline{\omega}_0(f) = 0$ if $\omega_0(f) \geq 0$ and $\overline{\omega}_0(f) = 1$ if $\omega_0(f) \leq -1$ and we set

$$\overline{N}(r, f) = \overline{\omega}_0(f) \log r + \sum_{n=1}^{\tau(r)} (\log r - \log |b_n|)$$
 and then $\overline{N}(r, f)$ is called *the counting function of the poles of f , ignoring multiplicity*

Now we can define the Nevanlinna function $T(r, f)$ in I or J as $T(r, f) = \max(Z(r, f), N(r, f))$ and the function $T(r, f)$ is called *characteristic function of f or Nevanlinna of f* .

Finally, if S is a subset of $\mathbb{K}$ we will denote by $Z_0^S(r, f')$ the counting function of zeros of f' , excluding those which are zeros of $f - a$ for any $a \in S$.

Remark: If we change the origin, the functions Z , N , T are not changed, up to an additive constant.

In a p -adic field such as $\mathbb{K}$, the first Main Theorem is almost immediate:

Theorem A: *Let $f \in \mathcal{M}(\mathbb{K})$ (resp. $f \in \mathcal{M}(d(0, R^-))$) have no zero and no pole at 0. Then $\log(|f|(r)) = \Psi(f, \log r) = \log(|f(0)|) + Z(r, f) - N(r, f)$.*

Corollary A.1: *Let $f, g \in \mathcal{M}(\mathbb{K})$ (resp. $f, g \in \mathcal{M}(d(0, R^-))$). Then $Z(r, fg) \leq Z(r, f) + Z(r, g)$, $N(r, fg) \leq N(r, f) + N(r, g)$, $T(r, fg) \leq T(r, f) + T(r, g)$, $T(r, f+g) \leq T(r, f) + T(r, g) + O(1)$, $T(r, cf) = T(r, f) \ \forall c \in \mathbb{K}^*$, $T(r, \frac{1}{f}) = T(r, f)$.*

If $f, g \in \mathcal{A}(\mathbb{K})$ (resp. $f, g \in \mathcal{A}(d(0, R^-))$), then $Z(r, fg) = Z(r, f) + Z(r, g)$, $T(r, f) = Z(r, f)$, $T(r, fg) = T(r, f) + T(r, g) + O(1)$ and $T(r, f+g) \leq \max(T(r, f), T(r, g))$.

We can now state the famous p -adic Second Main Theorem:

Theorem B: *Let $\alpha_1, \dots, \alpha_q \in \mathbb{K}$, with $q \geq 2$, let $S = \{\alpha_1, \dots, \alpha_q\}$ and let $f \in \mathcal{M}(\mathbb{K})$ (resp. $f \in \mathcal{M}_u(d(0, R^-))$). Then*

$$(q-1)T(r, f) \leq \sum_{j=1}^q \overline{Z}(r, f - \alpha_j) + \overline{N}(r, f) - Z_0^S(r, f') - \log r + O(1) \quad \forall r \in I \quad (\text{resp. } \forall r \in J).$$

Definitions and notation: For each $f \in \mathcal{M}(\mathbb{K})$ (resp. $f \in \mathcal{M}(d(a, R^-))$) we denote by $\mathcal{M}_f(\mathbb{K})$, (resp. $\mathcal{M}_f(d(a, R^-))$) the set of functions $h \in \mathcal{M}(\mathbb{K})$, (resp. $h \in \mathcal{M}(d(a, R^-))$) such that $T(r, h) = o(T(r, f))$ when r tends to $+\infty$ (resp. when r tends to R). Similarly,

if $f \in \mathcal{A}(\mathbb{K})$ (resp. $f \in \mathcal{A}(d(a, R^-))$) we shall denote by $\mathcal{A}_f(\mathbb{K})$ (resp. $\mathcal{A}_f(d(a, R^-))$) the set $\mathcal{M}_f(\mathbb{K}) \cap \mathcal{A}(\mathbb{K})$, (resp. $\mathcal{M}_f(d(a, R^-)) \cap \mathcal{A}(d(a, R^-))$).

The elements of $\mathcal{M}_f(\mathbb{K})$ (resp. $\mathcal{M}_f(d(a, R^-))$) are called *small functions with respect to f* . Similarly, if $f \in \mathcal{A}(\mathbb{K})$ (resp. $f \in \mathcal{A}(d(a, R^-))$) the elements of $\mathcal{A}_f(\mathbb{K})$ (resp. $\mathcal{A}_f(d(a, R^-))$) are called *small functions with respect to f* .

According to classical results [11], [18], we have the following Theorem C:

Theorem C: *Let $f \in \mathcal{A}(\mathbb{K})$ (resp. $f \in \mathcal{A}_u(d(0, R^-))$) and let $w_1, w_2 \in \mathcal{A}_f(\mathbb{K})$ (resp. $w_1, w_2 \in \mathcal{A}_f(d(0, R^-))$) be distinct. Then $T(r, f) \leq \bar{Z}(r, f - w_1) + \bar{Z}(r, f - w_2) + o(T(r, f))$.*

Definitions and notation: Given $f, g \in \mathcal{M}(d(0, R^-))$, we denote by $W(f, g)$ the Wronskian of f and g i.e. $f'g - fg'$.

In [5], the following results are proven:

Theorem D: *Let $f, g \in \mathcal{A}(\mathbb{K})$ be such that $W(f, g)$ is a non-identically zero polynomial. Then both f, g are polynomials.*

Notation: Let $f \in \mathcal{A}(\mathbb{K})$. We can factorize f in the form $\bar{f}\tilde{f}$ where the zeros of $\bar{f}$ are the distinct zeros of f each with order 1. Moreover, if $f(0) \neq 0$ we will take $\bar{f}$ such that $\bar{f}(0) = 1$.

Theorem E: *Let $f \in \mathcal{M}(\mathbb{K})$ have finitely many multiple poles, such that for certain $b \in \mathbb{K}$, $f' - b$ has finitely many zeros. Then f belongs to $\mathbb{K}(x)$.*

Notation: Let $f \in \mathcal{M}(d(0, R^-))$. For each $r \in]0, R[$, we denote by $\zeta(r, f)$ the number of zeros of f in $d(0, r)$, taking multiplicity into account and set $\tau(r, f) = \zeta(r, \frac{1}{f})$. Similarly, we denote by $\beta(r, f)$ the number of multiple zeros of f in $d(0, r)$, each counted with its multiplicity and we set $\gamma(r, f) = \beta(r, \frac{1}{f})$.

Theorem F: *Let $f \in \mathcal{M}(\mathbb{K})$ be such that for some $c, d \in]0, +\infty[$, $\gamma(r, f)$ satisfies $\gamma(r, f) \leq cr^d$ in $[1, +\infty[$. If f' has finitely many zeros, then $f \in \mathbb{K}(x)$.*

Corollary F.1: *Let f be a meromorphic function on $\mathbb{K}$ such that, for some $c, d \in]0, +\infty[$, $\gamma(r, f)$ satisfies $\gamma(r, f) \leq cr^d$ in $[1, +\infty[$. If for some $b \in \mathbb{K}$ $f' - b$ has finitely many zeros, then f is a rational function.*

First results [17]

We will now prove together the following Theorems 1 and 2.

Theorem 1: Let $f \in \mathcal{M}(\mathbb{K}) \setminus \mathbb{K}(x)$ satisfy $\limsup_{r \rightarrow \infty} |f|(r) > 0$ and let $b \in \mathbb{K}^*$. Let $m \in \mathbb{N}^*$ be ≥ 3 . Then $f' + bf^m$ has infinitely many zeros that are not zeros of f .

Theorem 2: Let $f \in \mathcal{M}_u(d(a, R^-))$ satisfy $\limsup_{r \rightarrow R} |f|(r) = +\infty$ and let $b \in \mathbb{K}^*$. Let $m \in \mathbb{N}^*$ be ≥ 3 . Then $f' + bf^m$ has infinitely many zeros that are not zeros of f .

Proof : Without loss of generality, we can assume $b = 1$ and when $f \in \mathcal{M}(d(a, R^-))$, we may assume $a = 0$. By hypotheses, there exists a sequence of intervals $[r'_n, r''_n]$ such that $\lim_{n \rightarrow +\infty} r'_n = +\infty$ (resp. $\lim_{n \rightarrow +\infty} r'_n = \lim_{n \rightarrow +\infty} r''_n = R$) and such that, putting $J = \bigcup_{n \in \mathbb{N}} [r'_n, r''_n]$, we have $\liminf_{\substack{r \rightarrow \infty, \\ r \in J}} |f|(r) > 0$ (resp. $\lim_{\substack{r \rightarrow R^-, \\ r \in J}} |f|(r) = +\infty$).

Suppose first we assume the hypothesis of Theorem 1. Let $M = \frac{\liminf_{r \rightarrow +\infty} |f|(r)}{2}$. We will prove that there exists $t > 0$ such that $|f' + f^m|(r) = |f^m|(r) \forall r \in J \cap [t, +\infty[$. We know that $|f'|(r) \leq \frac{|f|(r)}{r}$. Consequently, when r lies in J , there exists $s > 0$ such that $|f|(r) \geq M \forall r \in [s, +\infty[\cap J$.

$$(|f|(r))^m \geq |f|(r)M^{m-1} \geq r|f'|(r)M^{m-1}.$$

Next, when r is big enough, rM^{m-1} is greater than 1, hence $(|f|(r))^m > |f'|(r)$. Thus there exists $t \geq s$ such that $(|f|(r))^m > |f'|(r) \forall r \in J \cap [t, +\infty[$. Let $J' = J \cap [t, +\infty[$. And hence we have $|f' + f^m|(r) = |f^m|(r) \forall r \in J'$.

Suppose now that we assume the hypothesis of Theorem 2. We have $|f'|(r) \leq \frac{|f|(r)}{r} \leq \frac{|f|(r)}{R}$. Set $B = \frac{1}{R}$. Then we have

$$(|f|(r))^m \geq B|f'|(r)(|f|(r))^{m-1}.$$

Now, when r is close enough to R , $r \in J$, $B|f|(r)^{m-1}$ is strictly greater than 1, hence $(|f|(r))^m > |f'|(r)$. Thus there exists $t > 0$ such that $(|f|(r))^m > |f'|(r) \forall r \in [t, +\infty[\cap J$. We can set again $J' = J \cap [t, R[$ and then we have $|f' + f^m|(r) = |f^m|(r) \forall r \in J'$.

We can now conclude in both theorems 1 and 2. For each $n \in \mathbb{N}$, let q_n be the number of zeros of f in $d(0, r''_n)$. Suppose the sequence $(q_n)_{n \in \mathbb{N}}$ is bounded. Then, f has finitely many zeros, hence it is of the form $\frac{P}{h}$ with $P \in \mathbb{K}[x]$ and $h \in \mathcal{A}_u(d(0, R))$. Consequently, we have $\lim_{r \rightarrow +\infty} |f|(r) = 0$, a contradiction to the hypothesis in both theorems. Therefore, the sequence $(q_n)_{n \in \mathbb{N}}$ which is increasing by definition, tends to $+\infty$. Now, in each Theorems 1 and 2 we may apply Lemma 3 showing that $f' + f^m$ has infinitely many zeros that are not zeros of f .

Consider now the case $m = 1$. We can have a better conclusion in $\mathcal{M}(\mathbb{K})$.

Theorem 3: *Let $f \in \mathcal{M}(\mathbb{K}) \setminus \mathbb{K}(x)$. For each $b \in \mathbb{K}^*$, $f' + bf$ has infinitely many zeros that are not zeros of f .*

Proof: Without loss of generality, we can assume again $b = 1$. We have $|f'|(r) < |f|(r)$ when r is big enough and hence $|f' + f|(r) = |f|(r)$ in an interval $I = [s, +\infty[$. Suppose first that f has infinitely many zeros. We can then apply Lemma 3 and get the conclusion.

Suppose now that f has finitely many zeros. Then f has infinitely many poles c_n of respective order t_n . Since $\mathbb{K}$ has characteristic zero, f' admits each c_n as a pole of order $t_n + 1$ and similarly, $f' + f$ also admits each c_n as a pole of order $t_n + 1$. Thus, we have $N(r, f' + f) = N(r, f) + \overline{N}(r, f)$. But since $|f' + f|(r) = |f|(r)$ holds in I , we have $\Psi(f' + f, \log r) = \Psi(f, \log r) \forall r \in I$ hence, by Lemma 2, $\nu(f' + f, \log r) = \nu(f, \log r) \forall r \in I$ and hence $Z(r, f' + f) - N(r, f' + f) = Z(r, f) - N(r, f)$, therefore $Z(r, f' + f) - (N(r, f) + \overline{N}(r, f)) = Z(r, f) - N(r, f)$ and hence $Z(r, f' + f) = Z(r, f) + \overline{N}(r, f)$. Since we have supposed that f has finitely many zeros and since f has infinitely many poles, $f' + f$ has infinitely many zeros and all but finitely many are not zeros of f .

Theorem 4: *Let $f \in \mathcal{A}(\mathbb{K}) \setminus \mathbb{K}(x)$ (resp. let $a \in \mathbb{K}$, let $R \in]0, +\infty[$ and let $f \in \mathcal{A}_u(da, (R^+))$). For each $b \in \mathbb{K}^*$, $f' + bf^2$ has infinitely many zeros that are not zeros of f .*

Proof: Without loss of generality, we can assume $b = 1$ and $a = 0$. Clearly, when r is big enough, in $]0, +\infty[$ (resp. in $]0, R[$), we have $|f' + f^2|(r) = |f^2|(r)$ therefore f^2 and $f' + f^2$ have the same number of zeros in $C(0, r)$. Let $\alpha \in C(0, r)$ be a zero of f of order q . When r is big enough, it is a zero of order $2q$ for f^2 and it is a zero of order $q - 1$ for $f' + f^2$. Consequently, $f' + f^2$ has at least $q + 1$ zero in $C(0, r)$ that are not zeros of f (taking multiplicity into account). This is true for every such zeros of f and hence $f' + f^2$ has infinitely many zeros that are not zeros of f .

Corollary 4.1: *Let $m \in \mathbb{N}$ be ≥ 1 , let $f \in \mathcal{A}(\mathbb{K}) \setminus \mathbb{K}(x)$. For each $b \in \mathbb{K}^*$, $f' + bf^m$ has infinitely many zeros that are not zeros of f .*

Corollary 4.2: *Let $m \in \mathbb{N}$ be ≥ 2 , let $a \in \mathbb{K}$, let $R \in]0, +\infty[$ and let $f \in \mathcal{A}_u(da, (R^-))$. For each $b \in \mathbb{K}^*$, $f' + bf^m$ has infinitely many zeros that are not zeros of f .*

Theorem 5 is given in [4]:

Theorem 5: *Let $f \in \mathcal{M}(\mathbb{K}) \setminus \mathbb{K}(x)$ (resp. let $a \in \mathbb{K}$ and $R \in \mathbb{R}_+^*$ and let $f \in \mathcal{M}(d(a, R^-))$) and let $m \in \mathbb{N}$. If $m \geq 5$ then for each $b \in \mathbb{K}^*$, $f' + bf^m$ has infinitely many zeros that are not zeros of f .*

If $m = 4$, if $f \in \mathcal{M}(\mathbb{K}) \setminus \mathbb{K}(x)$ and if f admits at least s multiple zeros and at least t multiple poles, then $f' + bf^4$ admits a number of zeros that are not zeros of f (taken account of multiplicity) which is strictly superior to $\frac{s+t}{2}$.

Proof: We know that the zeros of $f' + bf^m$ in $\mathbb{K}$ are the same as in a spherically complete algebraically closed extension $\widehat{\mathbb{K}}$ of $\mathbb{K}$. So, for simplicity, we can suppose that the field $\mathbb{K}$

is spherically complete without loss of generality. We can also suppose that $b = 1$. Then if $f \in \mathcal{M}(\mathbb{K}) \setminus \mathbb{K}(x)$ we can obviously write $f = \frac{h}{l}$ with $h, l \in \mathcal{A}(\mathbb{K})$, having no common zeros and if $f \in \mathcal{M}(d(a, R^-))$, since $\mathbb{K}$ is spherically complete, we can write $f = \frac{h}{l}$ with $h, l \in \mathcal{A}(d(a, R^-))$, having no common zeros again.

Let $g = \frac{1}{f}$ and let $n = m - 2$. So, by Lemma 1, the problem is reduced to show that

$g'g^n - 1$ has infinitely many zeros. Then, $g'g^n - 1 = \frac{(l'h - hl')l^n - h^{n+2}}{h^{n+2}}$ and since h, l

have no common zeros, this is of the form $\frac{P}{h^{n+2}}$ where P is a polynomial of degree q .

Now, set $F = (l'h - hl')l^n$. Applying Theorem C to F we have

(1) $T(r, F) = Z(r, F) + O(1) \leq \bar{Z}(r, F) + \bar{Z}(r, F - P) + T(r, P) + O(1)$. By (1) we derive $Z(r, l'h - hl') + nZ(r, l) \leq \bar{Z}(r, l'h - hl') + \bar{Z}(r, l) + \bar{Z}(r, F - P) + T(r, P) + O(1)$. Actually, $\bar{Z}(r, F - P) = \bar{Z}(r, h)$, hence $nZ(r, l) \leq \bar{Z}(r, l) + \bar{Z}(r, h) + T(r, P) + O(1)$ and hence $(n - 1)Z(r, l) \leq Z(r, h) + T(P) + O(1)$. But since $T(r, P) = q \log r + O(1)$, we have

(2) $(n - 1)Z(r, l) \leq Z(r, h) + q \log r + O(1)$

Now, consider the hypothesis $f \in \mathcal{M}(\mathbb{K})$. By Theorem 1, if $\liminf_{r \rightarrow +\infty} |f|(r) > 0$ i.e. if $\liminf_{r \rightarrow +\infty} Z(r, f) - N(r, f) > -\infty$ the claim is proved. Consequently, if the claim is not true, we can assume

(3) $\liminf_{r \rightarrow +\infty} Z(r, f) - N(r, f) = -\infty$

But we see that (3) is impossible whenever $n \geq 3$, i.e. $m \geq 5$.

Now, suppose $m = 4$ i.e. $n = 2$. More precisely $\bar{Z}(r, l) \leq Z(r, l) - \frac{s \log r}{2}$ and $\bar{Z}(r, h) \leq Z(r, h) - \frac{t \log r}{2}$, so by Relation (1) we have

(4) $(n - 1)Z(r, l) \leq Z(r, h) + (q - \frac{s + t}{2}) \log r + O(1)$.

Then Relation (3) implies $q - \frac{s + t}{2} > 0$ and hence $f'f^n$ admits a number of zeros strictly superior to $\frac{s + t}{2}$.

Now, suppose that $f \in \mathcal{M}(d(0, R^-))$. By Theorem 2, if $\lim_{r \rightarrow R^-} |f|(r) = +\infty$ i.e. if $\liminf_{r \rightarrow R^-} Z(r, f) - N(r, f) = +\infty$ the claim is proved. Consequently, if the claim is not true, we can assume

(5) $\liminf_{r \rightarrow R^-} Z(r, f) - N(r, f) < +\infty$.

But by (2), we see that (5) is impossible whenever $n \geq 3$ i.e. $m \geq 5$.

Corollary 4.1: *Let $f \in \mathcal{M}(\mathbb{K}) \setminus \mathbb{K}(x)$. Then for each $n \geq 3$ $f'f^n$ has infinitely many zeros that are not zeros of f .*

Corollary 4.2: *Let $f \in \mathcal{M}(\mathbb{K}) \setminus \mathbb{K}(x)$ have s multiple zeros and t multiple poles. Let $b \in \mathbb{K}^*$. Then $f' + bf^4$ has at least $s+t+1$ zeros taking account of multiplicity. Particularly, if f has infinitely many multiple zeros or poles, then $f' + bf^4$ has infinitely many zeros that are not zeros of f .*

Corollary 4.3: *Let $f \in \mathcal{M}(\mathbb{K}) \setminus \mathbb{K}(x)$ have s multiple zeros and t multiple poles. Given $b \in \mathbb{K}^*$, If f has infinitely many multiple zeros or poles, then $f'f^2 - b$ has infinitely many zeros.*

Case $n = 2, m = 4$

We will now thoroughly examine the situation when $m = 4$ i.e. $n = 2$, as made in [9]. This requires several basic lemmas.

Lemma 4: *Let $f \in \mathcal{M}(\mathbb{K})$ be transcendental and such that f' has finitely many multiple zeros. Then $\frac{f''f}{(f')^2}$ has no quasi-exceptional value.*

Proof: Let $g = \frac{f}{f'}$. A pole of g is a zero of f' , hence by hypothesis, g has finitely many multiple poles. Consequently, by Theorem E, g' has no quasi-exceptional value. And hence neither has $1 - g'$. But $g' = \frac{(f')^2 - f''f}{(f')^2} = 1 - \frac{f''f}{(f')^2}$. Therefore $\frac{f''f}{(f')^2}$ has no quasi-exceptional value.

Lemma 5: *Let $f \in \mathcal{M}(\mathbb{K})$ be transcendental and have finitely many multiple zeros. Then $f''f + 2(f')^2$ has infinitely many zeros that are not zeros of f .*

Proof: Suppose first that f' has infinitely many multiple zeros. Since f has finitely many multiple zeros, the zeros of f' are not zeros of f except at most finitely many. Hence f' has infinitely many multiple zeros that are not zeros of f . And then, they are zeros of f'' , hence of $f''f + 2(f')^2$, which proves the statement.

So we are now led to assume that f' has finitely many multiple zeros. By Lemma 4 $\frac{f''f + 2(f')^2}{(f')^2}$ has infinitely many zeros. Let $c \in \mathbb{K}$ be a pole of order q of f . Without loss of generality, we can suppose $c = 0$. The beginning of the Laurent developpement of f at 0 is of the form $\frac{a_{-q}}{x^q} + \frac{\varphi(x)}{x^{q-1}}$ whereas $\varphi \in \mathcal{M}(\mathbb{K})$ has no pole at 0. Consequently, $\frac{f''f + 2(f')^2}{(f')^2}$ is of the form

$$\frac{(a_{-q})^2(3q^2 + q) + x\phi(x)}{(a_{-q})^2(q^2) + x\psi(x)}$$

whereas $\phi, \psi \in \mathcal{M}(\mathbb{K})$ have no pole at 0. So, the function $\frac{f''f + 2(f')^2}{(f')^2}$ has no zero at 0.

Therefore, each zero of $\frac{f''f + 2(f')^2}{(f')^2}$ is a zero of $f''f + 2(f')^2$ and hence $f''f + 2(f')^2$ has infinitely many zeros.

Now, let us show that the zeros of $f''f + 2(f')^2$ are not zeros of f , except maybe finitely many. Let c be a zero of $f''f + 2(f')^2$ and suppose that c is a zero of f . Then, it is a zero of f' and hence it is a multiple zero of f . But by hypotheses, f has finitely many multiple zeros, hence the zeros of $f''f + 2(f')^2$ are not zeros of f , except at most finitely many. That finishes proving the claim.

Lemma 6: *Let $f \in \mathcal{M}(\mathbb{K})$ be transcendental and let $b \in \mathbb{K}^*$ be such that $f^2f' - b$ has finitely many zeros. Then, $N(r, f) \leq Z(r, f) + O(1)$.*

Proof: Let $F = f^2f'$. Since $F - b$ is transcendental and has finitely many zeros, it is of the form $\frac{P(x)}{h(x)}$ with $h \in \mathcal{A}(\mathbb{K}) \setminus \mathbb{K}[x]$. Consequently, $|F|(r)$ is a constant when r is big enough and therefore, by Theorem A we have $Z(r, F) = N(r, F) + O(1)$ when r is big enough. Now, $Z(r, F) = 2Z(r, f) + Z(r, f')$ and, by Theorem A $Z(r, f') \leq Z(r, f) + \bar{N}(r, f) - \log r + O(1)$. On the other hand, by Theorem A again, we have $N(r, F) = 3N(r, f) + \bar{N}(r, f)$. Consequently, $3N(r, f) + \bar{N}(r, f) \leq 3Z(r, f) + \bar{N}(r, f) - \log r + O(1)$, which proves the claim.

Theorem 6: *Let $f \in \mathcal{M}(\mathbb{K}) \setminus \mathbb{K}(x)$ Then for each $b \in \mathbb{K}^*$, $f'f^2 - b$ has infinitely many zeros.*

Proof: Let $b \in \mathbb{K}$ and suppose that the claim is wrong, i.e. $f'f^2 - b$ has q zeros, taking multiplicity into account. By Theorem 5, we may assume that f has finitely many multiple zeros and finitely multiple poles. Set $F = f^2f'$. Then $F' = f(f''f + 2(f')^2)$. By Lemma 5, $f''f + 2(f')^2$ has infinitely many zeros that are not zeros of f . Consequently, F' admits for zeros: the zeros of f and the zeros of $f''f + 2(f')^2$. And by Lemma 4, there exists a sequence of zeros of $f''f + 2(f')^2$ that are not zeros of f .

Let $S = \{0, b\}$ and let $Z_0^S(r, F')$ be the counting function of zeros of F' when $F(x)$ is different from 0 and b . Since $F - b$ has finitely many zeros, the zeros c of F' which are not zeros of f cannot satisfy $F(c) = b$ except at most finitely many. Consequently, there are infinitely many zeros of F' counted by the counting function $Z_0^S(r, F')$ and hence for every fixed integer $t \in \mathbb{N}$, we have

$$(1) \quad Z_0^S(r, F') \geq t \log r + O(1).$$

Let us apply Theorem B to F . We have

$$(2) \quad T(r, F) \leq \bar{Z}(r, F) + \bar{Z}(r, F - b) + \bar{N}(r, F) - Z_0^S(r, F') - \log(r) + O(1).$$

Now, we have

$$(3) \quad \bar{Z}(r, F) \leq Z(r, f) + Z(r, f')$$

$$(4) \quad \overline{N}(r, F) = \overline{N}(r, f)$$

and since the number of zeros of $F - b$ is q , taking multiplicity into account,

$$(5) \quad \overline{Z}(r, F - b) \leq q \log r + O(1).$$

Consequently, by (2), (3), (4), (5) we obtain

$$(6) \quad T(r, F) \leq Z(r, f) + Z(r, f') + \overline{N}(r, f) - Z_0^S(r, F') + (q - 1) \log r + O(1).$$

On the other hand, by construction, $T(r, F) \geq Z(r, F) = 2Z(r, f) + Z(r, f')$ hence by (6) we obtain (7):

$$(7) \quad Z(r, f) \leq \overline{N}(r, f) - Z_0^S(r, F') + (q - 1) \log r + O(1).$$

Now, by Lemma 6, we have $N(r, f) \leq Z(r, f) + O(1)$ hence by (7) we obtain $0 \leq (q - 1) \log r - Z_0^S(r, F') + O(1)$ and hence by (1), fixing $t > q - 1$ we can derive $0 \leq (q - 1) \log r - t \log r + O(1)$, a contradiction. That finishes the proof of Theorem 6.

By Lemma 1, Theorems 5 and 6 we can now state the general result on the p -adic Hayman conjecture:

Corollary 6.1: *Let $f \in \mathcal{M}(\mathbb{K})$ be transcendental. Then for every $n \in \mathbb{N}$, $n \geq 2$, for every $b \in \mathbb{K}^*$, $f^2 f' - b$ has infinitely many zeros and for every $m \geq 4$, $f' + b f^m$ has infinitely many zeros that are not zeros of f .*

Case $n = 1$, $m = 3$

Concerning the case $m = 3$ i.e. $n = 1$ which remains unsolved, Corollary 6.1 has an immediate application to the conjecture with additional hypotheses [2].

Theorem 7: *Let $f \in \mathcal{M}(\mathbb{K})$. Suppose that there exists $c, d \in]0, +\infty[$, such that $\tau(r, f) \leq c r^d \forall r \in [1, +\infty[$. If $f' f^n - b$ has finitely many zeros for some $b \in \mathbb{K}$, with $n \in \mathbb{N}$, then $f \in \mathbb{K}(x)$.*

Proof: Suppose f is transcendental. By hypothesis, f^{n+1} satisfies $\zeta(r, \frac{1}{f^{n+1}}) = \tau(r, f^{n+1}) \leq c(n+1)r^d \forall r \in [1, +\infty[$ hence by Corollary 6.1, $f' f^n$ has no quasi-exceptional value.

Theorem 7 may be written in another way:

Corollary 7.1: *Let $f \in \mathcal{M}(\mathbb{K}) \setminus \mathbb{K}(x)$. Suppose that there exists $c, d \in]0, +\infty[$, such that $\zeta(r, f) \leq c r^d \forall r \in [1, +\infty[$. Then for all $m \in \mathbb{N}$, $m \geq 3$ and for all $b \in \mathbb{K}$, $f' - b f^m$ admits infinitely many zeros that are not zeros of f .*

Proof: We set $g = \frac{1}{f}$. Then by Theorem 7 $g'g^{m-2}$ has no quasi-exceptional value. Consequently, given $b \in \mathbb{K}^*$, $g'g^{m-2} + b$ has infinitely many zeros and hence $f' - bf^m$ has infinitely many zeros that are not zeros of f . Next, if $b = 0$, by Theorem F, f' has infinitely many zeros.

Theorem 8: Let $f \in \mathcal{M}(\mathbb{K})$. Suppose that there exists $c, d \in]0, +\infty[$, such that $\beta(r, f) \leq cr^d \forall r \in [1, +\infty[$. Then, for all $b \in \mathbb{K}$, $\frac{f'}{f^2} - b$ has infinitely many zeros.

Proof: Set $g = \frac{1}{f}$ again. Since the poles of g are the zeros of f , we have $\gamma(r, g) \leq cr^d$. Consequently, by Corollary F.1, g' has no quasi-exceptional value.

Remark: Using Theorem 8 to study the zeros of $f' - bf^2$ that are not zeros of f is not so immediate, as we will see below because of residues of f at poles of order 1. Of course, if $\frac{1}{f}$ is an affine function, $f' + f^2$ has no zeros, except if it is identically zero. And if it is not identically zero, the residue at the pole is not 1 in the general case.

Case $n = 0$ i.e. $m = 2$

As we noticed at the beginning, due to the counter-example provided by the function $\tan$, the case $n = 0$ has no solution in $\mathbb{C}$. However, we can notice certain conclusions.

Lemma 7: Let $f = \frac{h}{l} \in \mathcal{M}(\mathbb{K})$ with $h, l \in \mathcal{A}(\mathbb{K})$ having no common zero, let $b \in \mathbb{K}^*$ and let $a \in \mathbb{K}$ be a zero of $h'l - hl' + bh^2$ that is not a zero of $f' + bf^2$. Then a is a pole of order 1 of f and $\text{res}(f, a) = \frac{1}{b}$.

Proof: Clearly, if $l(a) \neq 0$, a is a zero of $f' + bf^2$. Hence, a zero a of $h'l - hl' + bh^2$ that is not a zero of $f' + bf^2$ is a pole of f . Now, when $l(a) = 0$, we have $h(a) \neq 0$ hence $l'(a) = bh(a) \neq 0$ and therefore a is a pole of order 1 of f such that $\frac{h(a)}{l'(a)} = \frac{1}{b}$. But since a is a pole of order 1, we have $\text{res}(f, a) = \frac{h(a)}{l'(a)}$, which ends the proof.

Theorem 9 is not a result specific to p -adic analysis but it will be useful in Theorem 10.

Theorem 9: Let $f \in \mathcal{M}(\mathbb{K})$, (resp. let $a \in \mathbb{K}$, let $f \in \mathcal{M}(d(a, R^-))$), let $b \in \mathbb{K}^*$ and let $\alpha \in \mathbb{K}$ (resp. let $\alpha \in d(a, R^-)$) be a point that is not a zero of f and such that the residue of f at α is different from $\frac{1}{b}$. Then α is a zero of $f' + bf^2$ if and only if it is a zero of $\frac{f'}{f^2} + b$. Moreover, if it is a zero of both functions, it has the same multiplicity with both.

Proof: Suppose first α is a zero of $f' + bf^2$. If α is not a pole of f , of course it is a zero of $\frac{f'}{f^2} + b$ with same multiplicity. Suppose now that α is a pole of f : since it is not a pole of $f' + bf^2$ it must be a pole of order 1 of f . Without loss of generality, we may assume that $\alpha = 0$ (resp. $a = \alpha = 0$). Consider the Laurent series of f at 0: $f(x) = \frac{a_{-1}}{x} + a_0 + a_1x + x^2\phi(x)$ with $\phi \in \mathcal{M}(\mathbb{K})$ (resp. $\phi \in \mathcal{M}(d(0, R^-))$) and $\phi(0) \neq \infty$. Then $f' + bf^2$ is of the form

$$f'(x) + bf(x)^2 = \frac{a_{-1}(-1 + ba_{-1})}{x^2} + \frac{2ba_0a_1}{x} + a_1 + b(a_0^2 + 2a_1a_{-1}) + x\xi(x)$$

with $\xi \in \mathcal{M}(\mathbb{K})$ (resp. $\xi \in \mathcal{M}(d(0, R^-))$) and $\xi(0) \neq \infty$ and hence, we have $a_{-1}(-1 + ba_{-1}) = 0$, $a_0a_{-1} = 0$, $a_0^2 + 2a_1a_{-1} = 0$. Since by hypothesis $\text{res}(f, \alpha) \neq -\frac{1}{b}$ we have $(1 + ba_{-1}) \neq 0$, hence $a_{-1} = 0$, a contradiction. Consequently, every zero of $f' + bf^2$ that is not a zero of f is a zero of $\frac{f'}{f^2} + b$ with same multiplicity.

Conversely, suppose now that α is a zero of $\frac{f'}{f^2} + b$. If α is not a pole of f , it is a zero of $f' + bf^2$, with the same multiplicity, because by hypothesis it is not a zero of f . Now suppose that α is a zero of $\frac{f'}{f^2} + b$ and is a pole of f . Clearly, it is a pole of order 1 and again, we may assume that $\alpha = 0$.

Consider again the Laurent series of f at 0: $f(x) = \frac{a_{-1}}{x} + a_0 + a_1x + x^2\phi(x)$ with $\phi \in \mathcal{M}(\mathbb{K})$ and $\phi(0) \neq \infty$. Then

$$\frac{f'}{f^2} = \frac{\frac{-a_{-1}}{x^2} + a_1 + x\psi(x)}{\frac{(a_{-1})^2}{x^2} + \frac{2a_0a_1}{x} + a_0^2 + 2a_1a_{-1} + x\xi(x)}$$

where both ψ , $\xi \in \mathcal{M}(\mathbb{K})$ have no pole at 0. Clearly, $\frac{f'}{f^2}$ is analytic at 0 and its value is $\frac{-1}{a_{-1}}$. But since 0 is a zero of $\frac{f'}{f^2} + b$, we have $a_{-1} = \frac{1}{b}$, what is excluded by hypothesis.

Thus we have proved that every zero of $\frac{f'}{f^2} + b$ is a zero of $f' + bf^2$ (that is not a zero of f) with the same multiplicity and this ends the proof of Theorem 9.

Theorem 10: Let $b \in \mathbb{K}^*$ and let $f \in \mathcal{M}(\mathbb{K})$ have finitely many zeros and finitely many residues at its simple poles equal to $\frac{1}{b}$ and be such that $f' + bf^2$ has finitely many zeros. Then f belongs to $\mathbb{K}(x)$.

Proof: Let $f = \frac{P}{l}$ with $P \in \mathbb{K}[x]$, $l \in \mathcal{A}(\mathbb{K})$ having no common zero with P . Then

$f' + bf^2 = \frac{P'l - l'P + bP^2}{l^2}$. By hypothesis, this function has finitely many zeros. Moreover, if a is a zero of $P'l - l'P + bP^2$ but is not a zero of $f' + bf^2$, then by Lemma 7 it is a pole of order 1 of f such that $\text{res}(f, a) = \frac{1}{b}$. Consequently, $P'l - l'P + bP^2$ has finitely many zeros and hence, we may write $\frac{P'l - l'P + bP^2}{l^2} = \frac{Q}{l^2}$ with $Q \in \mathbb{K}[x]$, hence $P'l - l'P = -bP^2 + Q$. But then, by Theorem D, l is a polynomial, which ends the proof.

Remark: If $f(x) = \frac{1}{x}$, the function $f' + bf^2$ has no zero whenever $b \neq 1$.

Theorem 11: *Let $f \in \mathcal{M}(\mathbb{K})$ be transcendental and have finitely many zeros of order ≥ 2 and let $b \in \mathbb{K}$. Then $\frac{f'}{f^2} + b$ has infinitely many zeros. Moreover, if $b \neq 0$, every zero α of $\frac{f'}{f^2} + b$ that is not a zero of $f' + bf^2$ is a pole of f of order 1 such that the residue of f at α is equal to $\frac{1}{b}$.*

Proof: Let $g = \frac{f'}{f^2} + b$. Since all zeros of f are of order 1 except maybe finitely many, g has finitely many poles of order ≥ 3 , hence a primitive G of g has finitely many poles of order ≥ 2 . Consequently, by Theorem E, g has infinitely many zeros.

Now, suppose $b \neq 0$. Let α be a zero of g . If α is not a pole of f , it is a zero of $f' + bf^2$ and we can see that it is not a zero of f .

Finally, suppose that α is a pole of f . Then it must be a pole of order 1 and then, by Lemma 7, the residue of f at α is $\frac{1}{b}$.

Corollary 11.1: *Let $f \in \mathcal{M}(\mathbb{K}) \setminus \mathbb{K}(x)$ have finitely many zeros of order ≥ 2 and finitely many poles of order 1 and let $b \in \mathbb{K}^*$. Then $f' + bf^2$ has infinitely many zeros that are not zeros of f .*

Remarks: As noticed above, in Archimedean analysis, the typical example of a meromorphic function f such that $f' - f^2$ has no zero is $\tan(x)$ and its residue is -1 at each pole of f . Here we find the same implication but we can't find an example satisfying such properties.

References

- [1] **Bergweiler, W. and Eremenko, A.** *On the singularities of the inverse to a meromorphic function of finite order*, Rev. Mat. Iberoamericana, 11, 355-373 (1995).
- [2] **Bezivin, J.-P., Boussaf, K., Escassut, A.** *Zeros of the derivative of a p -adic meromorphic function*, Bulletin des Sciences Mathématiques 136, 8, 839-847 (2012).

- [3] **Boussaf, K.** *Picard values of p -adic meromorphic functions*, *p-Adic Numbers Ultrametric Anal. Appl.* 2, no. 4, p.285-292 (2010).
- [4] **Boussaf, K. and Ojeda, J.** *Value distribution of p -adic meromorphic functions*, *Bull. Belg. Math. Soc. Simon Stevin*, 18, n.4, p. 667-678 (2011).
- [5] **Boussaf, K., Ojeda, J. and Escassut, A.** *Zeros of the derivative of a p -adic meromorphic function and applications* *Bull. Belg. Math. Soc. Simon Stevin* 19, n.2, p. 367-372 (2012).
- [6] **Boutabaa, A.** *Théorie de Nevanlinna p -adique*, *Manuscripta Math.* 67, p. 251-269 (1990).
- [7] **Boutabaa, A. and Escassut, A.,** *URS and URSIMS for p -adic meromorphic functions inside a disk*, *Proc. of the Edinburgh Mathematical Society* 44, p. 485-504 (2001).
- [8] **Escassut, A.** *Analytic Elements in p -adic Analysis*. World Scientific Publishing Co. Pte. Ltd. Singapore, (1995).
- [9] **Escassut, A. , Ojeda, J.** *The p -adic Hayman conjecture when $n = 2$* , *Complex variable and elliptic equations*.
- [10] **Hayman W. K.,** *Meromorphic Functions*. Oxford University Press, (1975)
- [11] **Hu, P.C. and Yang, C.C.** *Meromorphic Functions over non-Archimedean Fields*, Kluwer Academic Publishers, (2000).
- [12] **Hu, P.C. and Yang, C.C.** *Value Distribution Theory Related to Number Theory* Birkhäuser, (2006).
- [13] **Krasner, M.** *Prolongement analytique uniforme et multiforme dans les corps valués complets. Les tendances géométriques en algèbre et théorie des nombres*, Clermont-Ferrand, p.94-141 (1964). Centre National de la Recherche Scientifique (1966), (Colloques internationaux de C.N.R.S. Paris, 143).
- [14] **Lazard, M.** *Les zéros des fonctions analytiques sur un corps valué complet*, IHES, Publications Mathématiques no. 14, pp. 47-75 (1962).
- [15] **Mues, E.** *Über ein Problem von Hayman*, *Math. Z.*, 164, 239-259 (1979).
- [16] **Nevanlinna, R.** *Le théorème de Picard-Borel et la théorie des fonctions méromorphes*. Gauthiers-Villars, Paris (1929).
- [17] **Ojeda, J.** *On Hayman's Conjecture over a p -adic field*, *Taiwanese Journal of Mathematics* 12 (9), p. 2295-2313, (2008).
- [18] **Ojeda, J.** *Applications of the p -adic Nevanlinna theory to problems of uniqueness*, *Advances in p -adic and Non-Archimedean analysis. Contemporary Mathematics* 508, p.161-179 (2010).