Cost-effectiveness threshold of first-trimester Down syndrome maternal serum screening for the use of cell-free DNA as a second-tier screening test

Marie Blanquet, Sophie Dreux, Stéphanie Léger, Corinne Sault, Charline Mourgues, Hélène Laurichesse, Didier Lémery, Françoise Vendittelli, Anne Debost-Legrand, Françoise Muller

► To cite this version:

Marie Blanquet, Sophie Dreux, Stéphanie Léger, Corinne Sault, Charline Mourgues, et al.. Cost-effectiveness threshold of first-trimester Down syndrome maternal serum screening for the use of cell-free DNA as a second-tier screening test. *Annales de Biologie Clinique*, 2021, 79 (4), pp.331-338. 10.1684/abc.2021.1656 . hal-03439596

HAL Id: hal-03439596

<https://hal.uca.fr/hal-03439596>

Submitted on 22 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution| 4.0 International License

Titre : Evaluation de la valeur seuil la plus cout-efficace des marqueurs sériques du 1er trimestre pour l'emploi du test d'ADN circulant dans le dépistage de la trisomie 21 : une analyse médico-économique.

Titre court : Analyse cout-efficacité de la stratégie de dépistage de la trisomie 21

Title: Cost-effectiveness threshold of first-trimester Down syndrome maternal serum screening for the use of cell-free DNA as a second-tier screening test

Short-title: Cost-effectiveness study of Down syndrome screening strategy

Institute where the work was conducted: CHU of Clermont-Ferrand. France

Marie Blanquet^{1,2}, Sophie Dreux³, Stéphanie Léger⁴, Corinne Sault⁵, Charline Mourgues^{2,6},
Hélène Laurichesse^{2,7}, Didier Lémery^{2,7}, Françoise Vendittelli^{2,7}, Anne Debost-Legrand^{2,7},
Françoise Muller³.

¹ Service de Médecine, Centre Hospitalier de Mauriac, Mauriac, France

² Université Clermont Auvergne, CNRS-UMR 6602, Institut Pascal, Axe TGI, Péprade, Clermont-Ferrand, France

³ Laboratoire de Biochimie-Hormonologie, Hôpital Robert Debré, APHP, Paris, France

⁴ Laboratoire de mathématiques UMR CNRS 6620, Université Blaise Pascal; CNRS, UMR 6620, Laboratoire de mathématiques, Aubière, France.

⁵ Laboratoire Eurofins Biomnis, Lyon, France.

⁶ Direction de la Recherche Clinique, Centre Hospitalier Universitaire de Clermont-Ferrand, Clermont-Ferrand, France

⁷ Pôle Femme et Enfant, Centre Hospitalier Universitaire de Clermont-Ferrand, Clermont-Ferrand, France

1
2
3
4
5
6
7
8
9
10
11
12
13
14

Corresponding author

Dr Anne Debost-Legrand
CHU de Clermont-Ferrand, Pôle Femme et Enfant, place Lucie et Raymond Aubrac, 63003
Clermont-Ferrand
Phone number: 0033 473750715/Fax: 0033(0)473750619
Email address: alegrand@chu-clermontferrand.fr

Mots-clefs: test ADN libre circulant, Trisomie 21, dépistage prénatal, analyse économique,
ratio cout-efficacité,
Keywords: Non-invasive prenatal testing, Down syndrome, prenatal screening, economic
analysis, cost-effectiveness ratio.

Résumé

Introduction : Notre objectif était d'identifier la valeur seuil la plus cout-efficace des marqueurs sériques du 1^{er} trimestre pour l'intégration du test ADN libre circulant en deuxième ligne dans le cadre du dépistage prénatal de la trisomie 21 en France.

Matériel et méthodes : Une analyse cout-efficacité a été menée sur 108 121 grossesses uniques en utilisant un modèle de simulation. Les valeurs seuils testées des marqueurs sériques du 1^{er} trimestre étaient comprises entre 1/51 et 1/1000 par pas de 1/50. La valeur seuil la plus coût-efficace a été déterminée par le calcul du ratio coût-efficacité (RCE, coûts= coûts directs médicaux après le dosage des marqueurs du 1^{er} trimestre, efficacité=nombre de cas de trisomie 21 dépisté).

Résultat : Dans notre échantillon, 161 cas de trisomie 21 ont été identifiés. Pour une valeur seuil $\geq 1/50$, 47,2% des cas ont été diagnostiqués. Dans le modèle de simulation, pour un seuil $\geq 1/250$, 73,9% des cas de trisomie 21 étaient identifiés, pour un seuil $\geq 1/500$, le pourcentage de détection s'élevait à 78,8% et pour un seuil $\geq 1/1000$, 2 cas supplémentaires ont été identifiés. Le montant des coûts était stable pour un seuil $\geq 1/250$ (978 634€), puis augmentait rapidement pour un seuil $\geq 1/500$ (1 966 576€) et devenait exponentiel pour un seuil $\geq 1/1000$ (3 980 216€). Le RCE pour le seuil $\geq 1/500$ était de 38 560.

Conclusion : La valeur seuil la plus coût-efficace pour l'implémentation du test ADN libre circulant en deuxième ligne semble être $\geq 1/500$.

1 **Abstract**

2 Introduction: We aimed to identify the most relevant cost-effectiveness threshold of first-
3 trimester Down syndrome (DS) maternal serum screening (T21T1) for the use of cell-free
4 DNA (cfDNA) as a second-tier test in the French context.

5 Method: A cost-effectiveness analysis was performed on 108 121 singleton pregnancies
6 using a simulation model. The threshold of T21T1 screening was ranged from 1/51 to 1/1000
7 in steps of 1/50. The most relevant threshold was based on cost-effectiveness ratio (CER;
8 costs = direct medical costs after T21T1 screening/ effectiveness = number of DS cases
9 identified).

10 Results: In the sample, 161 cases of DS were identified. At the threshold of $\geq 1/50$, 47.2% of
11 total DS cases were diagnosed. In the simulation model, for a threshold $\geq 1/250$, 73.9% of
12 total DS cases were diagnosed, for $\geq 1/500$, 78.8% and for $\geq 1/1000$, only two additional
13 cases were diagnosed. The slope of the cost increase was slight from threshold $\geq 1/250$ (978
14 634€), then steep up to 1/500 (1 966 576€) and increased exponentially to 1/1000 (3 980
15 216€). The CER was 38 560 for a threshold $\geq 1/500$.

16 Conclusion: The most cost-effective threshold for cfDNA as a second-tier test seems to be $\geq$
17 1/500. For higher thresholds, costs increase dramatically for only a few additional cases of DS
18 identified.

19

Introduction

Over the last 50 years, prenatal diagnosis procedures and screening have become safer and the capabilities of diagnostic laboratories have expanded[1–6]. Recently, the use of cell-free DNA (cfDNA) has dramatically increased. While cfDNA is considered as a screening test and not a diagnostic test by the scientific community, patients and most physicians see cfDNA as a substitute for foetal karyotyping[7–9].

Recently, new guidelines drawn up by the French National Authority for Health (HAS; Haute Autorité de Santé) recommend cfDNA for Down syndrome (DS) screening[10]. These guidelines, based on a cost-effectiveness analysis, recommend cfDNA as a second-tier screening test in women with a risk between 1/51 and 1/1000 after first-trimester combined maternal serum screening (T21T1). Moreover, in high-risk women ($T21T1 \geq 1/50$), invasive testing is proposed but, after clear information, women still have the choice of cfDNA. However, these guidelines were based on simulated data.

CfDNA is a reliable technique that should be included in DS screening, but it remains contingent on maternal serum screening and the terms of application need to be carefully defined. It is necessary to identify the most relevant threshold of the calculated risk of DS based on maternal serum markers for the use of cfDNA. Few studies have tested different thresholds of T21T1 to identify the most cost-effective one[11–14]. Even though these studies reached similar conclusions, the number of thresholds tested was limited and there was no step-by-step analysis. The principal objective of this study was to identify the most relevant cost-effectiveness threshold of T21T1 screening for use of cfDNA as a second-tier test, by carrying out an economic analysis, in the French context.

Methods

French prenatal screening for Down syndrome

In France, prenatal DS screening has been regulated by the Ministry of Health since 1997[15]. The French DS screening policy is organized at a national scale and since 2010 has been based on three steps. The first consists of first-trimester nuchal translucency (NT) and crown rump length (CRL) measurements (T1US), a cut-off ≥ 3.5 mm leading to chorionic villus sampling (CVS) for karyotyping. The second step consists of first-trimester maternal serum screening (or second-trimester when NT is not performed or when maternal blood sampling is performed after 13⁺⁶ weeks of gestation). First-trimester DS maternal serum screening (T21T1) is performed at 11-13⁺⁶ weeks and combined maternal age, maternal serum markers (pregnancy-associated plasma protein A, PAPP-A, and free beta-human chorionic gonadotropin, hCG β) and nuchal translucency (NT). Karyotyping is offered for a calculated risk $\geq 1/50$. If women have a risk between 1/51 and 1/1000 after first-trimester combined maternal serum screening (T21T1), cfDNA is recommended as a second-tier screening test[10]. The third step consists of two ultrasound scan examinations at 20-22 weeks and 30-32 weeks of gestation.

Study population

Our economic analysis was applied to data from 108 121 singleton pregnancies with T21T1 (Autodelfia, dual kit and LifeCycle 4.1 software, PerkinElmer, Turku, Finland) during the year 2015 in the Biomnis Eurofins Laboratory, one of 86 accredited laboratories. Maternal serum markers and NT were expressed as multiple of the median (MoM) for gestational age (expressed in weeks and days of amenorrhea) as estimated by CRL measurement performed at the same time as NT. MoMs were adjusted for three confounding factors (maternal weight, smoking status and ethnicity). In accordance with French law, written informed consent for biochemical testing was obtained from each woman.

If chorionic villous sampling or amniocentesis was performed, a second written consent was needed for foetal karyotyping. The database has been reported to the French Data Protection Authority (CNIL: Commission Nationale de l'Informatique et des Libertés) as CNIL no. 1839545 V1.

Economic analysis

Identification of DS in the database

Cases in which embryo reduction was performed and cases in which a vanishing twin was detected at ultrasound scan were excluded. Because of the French policy using NT as contingent screening, all cases with $NT \geq 3.5$ mm were excluded from the analysis. DS cases were identified prenatally when women had T21T1 screening $\geq 1/250$ (based on French policy of DS screening strategy in 2015) or when anomalies were detected at ultrasound scan or at birth when the patient did not opt for an invasive procedure or for termination of pregnancy or when T21T1 was $< 1/250$ or without ultrasound anomalies. To sum up, among 108 121 singleton pregnancies 161 cases of DS were diagnosed, 124 by T21T1 screening, 16 by means of abnormal ultrasound scans and 21 at birth. All karyotyping results for the cohort were collected by the accredited laboratory, which performed serum marker screening.

Simulation model

The simulation model was based on actual data regarding DS status (including overall cases) and T21T1 results. Three variables were simulated based on assumptions from the international literature or expertise. Those variables were the women's characteristics (body weight and ethnicity), cfDNA (sensitivity, specificity and failure rate) and karyotyping acceptance rate. Women whose T21T1 results indicated a risk $\geq 1/1000$ were considered to be part of a high-risk group in the simulation model, so the acceptability of cfDNA testing and karyotyping was presumed to be high. Table 1 presents the percentages with the 95% confidence intervals of each variable included in the model.

Cost-effectiveness analysis

A cost-effectiveness analysis (CEA) was performed for different thresholds and calculated risk as a discrete variable of T21T1 screening was used to identify the most relevant threshold. Figure 1 depicted the decision tree model used in the analysis describing the clinical pathway where cfDNA was used as a second-tier test if T21T1 screening was positive starting from $\geq 1/50$ to $\geq 1/1000$ by step of $1/50$.

The perspective chosen was that of the third-party payer, ie, the French national health insurance fund (Caisse Nationale d'Assurance Maladie). Expressed in euros, the costs included were the direct medical costs after T21T1 screening: the cost of karyotyping, the invasive technique and the procedure used (chorionic villus sampling or amniocentesis) and of karyotype analysis (Table 1). Costs were calculated for the sample of 108 121 women. The incremental cost per DS case diagnosed was also calculated. The medical and societal costs of an undetected DS-affected child were not taken into account. The costs and effectiveness criteria were based on data from a single year and have not been actualized[16,17].

The most relevant threshold of T21T1 screening for use of cfDNA as a second-tier test was based on cost-effectiveness ratio (CER). The criterion of effectiveness considered was the number of DS cases diagnosed. The CER was calculated as follows: CER = total cost for each threshold/number of DS cases diagnosed at each threshold. Univariate sensitivity analysis was used to test the robustness of the results by varying the cost of cfDNA.

Statistical analysis

A descriptive analysis of the data base was done for quantitative variables with standard deviation and for qualitative variables with 95% confidence interval (95% CI). The cohort's advance through the decision tree was based on DS prevalence in the actual database, cfDNA (sensitivity and specificity) and screening test participation rates for karyotyping (Table 1)

[16–21]. The variation in the failure rate of the cfDNA test as a function of body weight was taken into consideration as was ethnicity[18,22]. Foetal loss due to invasive screening was also considered for each threshold[19]. The threshold of T21T1 screening ranged from 1/51 to 1/1000 in steps of 1/50. We used R 3.0.2. Software for this modelling.

Results

Descriptive analysis of actual data

A total of 108 121 women were included in the study. The mean maternal age was 29.7 [standard deviation 4.9] years. Median measurement of NT was 1.4 [95% CI: 0.8; 2.1]. Mean gestational age at T21T1 was 12^{+6} WG (11^{+0} - 13^{+4}). In the sample, DS was diagnosed in 161 cases, giving a 0.2% [95% CI: 0.1; 0.2] global prevalence.

At the threshold of $\geq 1/50$, among 76 cases of DS diagnosed, 71 were confirmed by foetal karyotyping and 5 after birth when the parents refused prenatal karyotyping (Table 2).

Results of the simulation model

For thresholds at 1/51-1/250, 43 additional cases were diagnosed (119 cases, 73.9% of total DS cases). Then, for thresholds at 1/251-1/500, 8 additional cases were diagnosed (127 cases, 78.9% of total DS cases). Finally, for thresholds at 1/501-1/1000, 2 additional cases were diagnosed (129 cases, 80.1% of total DS cases). One foetal loss was observed for a threshold of 1/251 (Table 2).

Considering that a cfDNA test costs 330 €, the slope of the cost increase was slight from a threshold 1/51-1/250 (978 634 €), then steep up to a threshold 1/251-1/500 (1 966 576 €) and increased exponentially from 1/501 to 1/1000 (3 980 215 €) (Figure 2). From a threshold 1/750 to 1/1000, no additional DS case was diagnosed, whereas the cost still increased (Figure 2). The CER was 38 560 for threshold 1/51-1/500, 56 471 at 1/51-1/750 and 75 098 at 1/51-1/1000. The incremental cost per DS case diagnosed was 191 637 € for threshold 1/51-1/500,

406 014 € at 1/51-1/750 and 735 092 € at 1/51-1/1000. The results were robust irrespective of the cost of cfDNA (Figure 2).

Consequently, the most cost-effective threshold seems to be 1/51-1/500 as 127 (79.9% of total DS cases) DS cases were identified for a total cost of 1 966 576 €, a CER of 38 560 and an incremental cost per DS diagnosed of 191 637 € for a cfDNA cost of 330 € (Table 3).

Discussion

In our study based on actual data, the most cost-effective threshold of cfDNA as a second-tier test was 1/51-1/500. For higher thresholds, costs increase dramatically for the diagnosis of only a few additional cases of DS.

This is the first economic study based on actual data in the general population to assess by a step by step analysis the inclusion of cfDNA in the French strategy of DS screening, following new national guidelines. The study was based on a large sample: 20.4% of women who underwent T21T1 in 2015[23]. Because of exclusion of high NT, the prevalence of DS in our sample is 0.2% whereas in France in 2012 the estimated prevalence was 0.3%[24]. However, even if the prevalence of DS in our sample is low, the percentage of DS cases identified by screening remains stable, especially at the threshold from 1/51-1/750 to 1/51-1/1000, where no more cases of DS were identified.

First-trimester screening and cfDNA (based on the 1/1000 threshold[10]) identified up to 129 cases (corresponding to a threshold 1/51-1/750) and missed 11 cases, all of which were diagnosed by prenatal karyotyping. Considering the most cost-effective threshold of 1/51-1/500, only two more DS cases were missed, but they may be identified later by second-trimester ultrasound scan as it has been demonstrated that more than 30% of fetuses with DS can be detected among women classified as low risk after T21T1 screening[25]. Different costs for the cfDNA test were considered in a univariate sensitivity analysis. One limitation of

our study is that non-medical indirect costs were not considered[26]. The modelling did not consider the parents' decision based on the screening results. The major ethical and legal principle governing screening and prenatal diagnosis in France is that the final decision to continue or terminate the pregnancy belongs to the parents[27,28]. Other chromosomal aberrations such as trisomy 13 or 18 and sex chromosome abnormalities can be detected by cfDNA[29], however only results for DS were considered as it is the only chromosomal anomaly implemented in the national screening programme.

In our study, only the cfDNA contingent strategy was considered since most studies have concluded that universal cfDNA is nowadays too expensive and so irrelevant[13,30]. We are also in accordance with current national French guidelines which recommend cfDNA as a second-tier test[10]. Few studies have tested several thresholds to identify the most cost-effective cfDNA contingent strategy[11,12,14]. Chitty et al., Morris et al. and Nyet et al. demonstrated that the cfDNA contingent strategy remained cost-effective until a threshold of $\geq 1/150$, $\geq 1/500$ and $\geq 1/600$, respectively[11,12,14]. This is in line with our results regarding the dramatic cost increase at $<1/500$. The marginal cost of the cfDNA contingent strategy is very sensitive to the cost of cfDNA. In our study, for high-risk women ($T21T1 \geq 1/50$) karyotyping was systematically proposed. To our knowledge, no study has implemented this strategy in its model.

Based on a large series of actual data, the most cost-effective threshold for cfDNA as a second-tier test in first-trimester screening seems to be $1/51$ - $1/500$. In the literature, the optimal threshold is between $\geq 1/150$ and $\geq 1/600$ according to current national guidelines of each country, assuming an unchanging budget[11,12,14,31]. For higher thresholds, costs increase dramatically for only a few additional cases of DS identified. In a recent meta-analysis of economic evaluations of prenatal screening by cfDNA testing, the authors demonstrated that despite heterogeneity between studies mainly because of several time-

horizon considerations and different outcomes, the contingent strategy provides the best value for money[30]. On the other hand, the universal strategy appears cost-effective when cost of cfDNA decreases. In our study, the cost of cfDNA is the one used by the French National Table of Laboratory Codes and has not been yet actualized. As already underlined, cost-effectiveness studies results depend mainly of the cost of cfDNA. Another difference is also that many studies with simulated data include controversial future costs for children born with disabilities. However, a lifetime perspective is not appropriate in evaluating the prenatal screening strategy, as it appears unethical to estimate the value of life [32]. In the same way, we did not evaluate the viability of each pregnancy, as our objective was to simulate the foetal loss induced by invasive procedures. Nonetheless, the psychosocial consequences for families of these test results can be considerable and should be evaluated in economic studies[33]. Our choice to evaluate the hospital, rather than societal, point of view of the costs generated by positive test results means that we could not assess the costs of their psychosocial consequences and their effects on pregnancy outcomes[34]. Although these intangible costs are difficult to quantify, they can be approached by questionnaires, which should be used to show the full value of the DS screening strategy to health policy decision makers.

We have demonstrated in accordance with the literature that the contingent strategy is cost-effective when cfDNA testing is performed at a threshold $\geq 1/500$. National guidelines should take into account these results, especially in terms of opportunity costs for French society. Sensitivity analyses may be performed to assess validity of the model when cost of cfDNA will decrease as it will be used in routine screening in many countries. New thresholds may be more accurate than those used. The new challenge is to ensure that health professionals have adequate prenatal screening understanding in order to protect women's autonomy when consenting to genetic testing.

1 **Conflict of interest**

2 The authors declare that they have no conflict of interest.

References

1. Evans M, Johnson M, Yaron Y, Drugan A. Prenatal diagnosis : genetics, reproductive risks, testing and management. McGraw Hill Publishing Co. New York; 2006.
2. Chiu RWK, Chan KCA, Gao Y, Lau VYM, Zheng W, Leung TY, et al. Noninvasive prenatal diagnosis of fetal chromosomal aneuploidy by massively parallel genomic sequencing of DNA in maternal plasma. *Proc Natl Acad Sci U S A*. 2008;105:20458–63.
3. Ehrich M, Deciu C, Zwiefelhofer T, Tynan JA, Cagasan L, Tim R, et al. Noninvasive detection of fetal trisomy 21 by sequencing of DNA in maternal blood: a study in a clinical setting. *Am J Obstet Gynecol*. 2011;204:205.e1-11.
4. Evans MI, Krantz DA, Hallahan TW, Galen RS. Meta-analysis of first trimester Down syndrome screening studies: free beta-human chorionic gonadotropin significantly outperforms intact human chorionic gonadotropin in a multimarker protocol. *Am J Obstet Gynecol*. 2007;196:198–205.
5. Bianchi DW, Simpson JL, Jackson LG, Elias S, Holzgreve W, Evans MI, et al. Fetal gender and aneuploidy detection using fetal cells in maternal blood: analysis of NIFTY I data. National Institute of Child Health and Development Fetal Cell Isolation Study. *Prenat Diagn*. 2002;22:609–15.
6. Wapner RJ, Martin CL, Levy B, Ballif BC, Eng CM, Zachary JM, et al. Chromosomal microarray versus karyotyping for prenatal diagnosis. *N Engl J Med*. 2012;367:2175–84.
7. Benn P, Chapman AR, Erickson K, Defrancesco MS, Wilkins-Haug L, Egan JFX, et al. Obstetricians and gynecologists' practice and opinions of expanded carrier testing and noninvasive prenatal testing. *Prenat Diagn*. 2014;34:145–52.
8. Brewer J, Demers L, Musci T. Survey of US obstetrician opinions regarding NIPT use in general practice: implementation and barriers. *J Matern-Fetal Neonatal Med Off J Eur Assoc Perinat Med Fed Asia Ocean Perinat Soc Int Soc Perinat Obstet*. 2017;30:1793–6.
9. Chan WV, Johnson J-A, Wilson RD, Metcalfe A. Obstetrical provider knowledge and attitudes towards cell-free DNA screening: results of a cross-sectional national survey. *BMC Pregnancy Childbirth*. 2018;18:40.
10. Haute Autorité de santé. Place des tests ADN libre circulant dans le sang maternel dans le dépistage de la trisomie 21 foetale [Internet]. 2017 [cited 2018 Jun 12]. Available from: https://www.has-sante.fr/portail/jcms/c_2768510/fr
11. Neyt M, Hulstaert F, Gyselaers W. Introducing the non-invasive prenatal test for trisomy 21 in Belgium: a cost-consequences analysis. *BMJ Open* [Internet]. 2014 [cited 2015 Sep 4];4. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4225226/>
12. Morris S, Karlsen S, Chung N, Hill M, Chitty LS. Model-Based Analysis of Costs and Outcomes of Non-Invasive Prenatal Testing for Down's Syndrome Using Cell Free Fetal DNA in the UK National Health Service. *PLoS ONE* [Internet]. 2014 [cited 2015 Sep 4];9. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3979704/>

- 1 13. García-Pérez L, Linertová R, Álvarez-de-la-Rosa M, Bayón JC, Imaz-Iglesia I, Ferrer-
2 Rodríguez J, et al. Cost-effectiveness of cell-free DNA in maternal blood testing for prenatal
3 detection of trisomy 21, 18 and 13: a systematic review. *Eur J Health Econ HEPAC Health*
4 *Econ Prev Care*. 2017;
- 5 14. Chitty LS, Wright D, Hill M, Verhoef TI, Daley R, Lewis C, et al. Uptake, outcomes, and
6 costs of implementing non-invasive prenatal testing for Down's syndrome into NHS
7 maternity care: prospective cohort study in eight diverse maternity units. *BMJ*.
8 2016;354:i3426.
- 9 15. Arrêté du 23 juin 2009 fixant les règles de bonnes pratiques en matière de dépistage et de
10 diagnostic prénatals avec utilisation des marqueurs sériques maternels de la trisomie 21.
- 11 16. Caisse Nationale d'Assurance Maladie. Cotation des Actes Médicaux version 39.10
12 applicable au 01 janvier 2015. Caisse Nationale d'Assurance Maladie; 2014 p. 403.
- 13 17. Caisse Nationale d'Assurance Maladie. Table Nationale de codage de Biologie. Caisse
14 Nationale d'Assurance Maladie; 2015 Nov.
- 15 18. Ashoor G, Syngelaki A, Poon LCY, Rezende JC, Nicolaides KH. Fetal fraction in
16 maternal plasma cell-free DNA at 11-13 weeks' gestation: relation to maternal and fetal
17 characteristics. *Ultrasound Obstet Gynecol Off J Int Soc Ultrasound Obstet Gynecol*.
18 2013;41:26–32.
- 19 19. Akolekar R, Beta J, Picciarelli G, Ogilvie C, D'Antonio F. Procedure-related risk of
20 miscarriage following amniocentesis and chorionic villus sampling: a systematic review and
21 meta-analysis. *Ultrasound Obstet Gynecol Off J Int Soc Ultrasound Obstet Gynecol*.
22 2015;45:16–26.
- 23 20. Gil MM, Quezada MS, Revello R, Akolekar R, Nicolaides KH. Analysis of cell-free DNA
24 in maternal blood in screening for fetal aneuploidies: updated meta-analysis. *Ultrasound*
25 *Obstet Gynecol Off J Int Soc Ultrasound Obstet Gynecol*. 2015;45:249–66.
- 26 21. Cuckle H, Benn P, Pergament E. Maternal cfDNA screening for Down syndrome--a cost
27 sensitivity analysis. *Prenat Diagn*. 2013;33:636–42.
- 28 22. Tanguy J, Zeghnoun A, Dor F. Description du poids corporel en fonction du sexe et de
29 l'âge dans la population française. *Environ Risques Santé*. 2007;6:179–87.
- 30 23. Agence de la biomédecine - rapport médical et scientifique [Internet]. [cited 2018 Feb 8].
31 Available from: [https://www.agence-biomedecine.fr/annexes/bilan2016/donnees/diag-](https://www.agence-biomedecine.fr/annexes/bilan2016/donnees/diag-prenat/01-diag_prenat/synthese.htm)
32 [prenat/01-diag_prenat/synthese.htm](https://www.agence-biomedecine.fr/annexes/bilan2016/donnees/diag-prenat/01-diag_prenat/synthese.htm)
- 33 24. Institut de Veille Sanitaire. Malformations congénitales et anomalies chromosomiques
34 [Internet]. [cited 2018 May 17]. Available from: [http://invs.santepubliquefrance.fr/Dossiers-](http://invs.santepubliquefrance.fr/Dossiers-thematiques/Maladies-chroniques-et-traumatismes/Malformations-congenitales-et-anomalies-chromosomiques/Donnees)
35 [thematiques/Maladies-chroniques-et-traumatismes/Malformations-congenitales-et-anomalies-](http://invs.santepubliquefrance.fr/Dossiers-thematiques/Maladies-chroniques-et-traumatismes/Malformations-congenitales-et-anomalies-chromosomiques/Donnees)
36 [chromosomiques/Donnees](http://invs.santepubliquefrance.fr/Dossiers-thematiques/Maladies-chroniques-et-traumatismes/Malformations-congenitales-et-anomalies-chromosomiques/Donnees)
- 37 25. Bromley B, Shipp TD, Lyons J, Groszmann Y, Navathe RS, Benacerraf BR. What is the
38 importance of second-trimester “soft markers” for trisomy 21 after an 11- to 14-week
39 aneuploidy screening scan? *J Ultrasound Med Off J Am Inst Ultrasound Med*. 2014;33:1747–
40 52.

26. Verhoef TI, Daley R, Vallejo-Torres L, Chitty LS, Morris S. Time and travel costs incurred by women attending antenatal tests: A costing study. *Midwifery*. 2016;40:148–52.
27. Comité Consultatif National d’Ethique. Avis sur les problèmes posés par le diagnostic prénatal et périnatal. Comité consultatif national d’Ethique; 1985 May p. 14. Report No.: 005.
28. Code de la Santé Publique. LOI n° 2011-814 du 7 juillet 2011 relative à la bioéthique. 2011-814 Jul 7, 2011.
29. Benachi A, Letourneau A, Kleinfinger P, Senat M-V, Gautier E, Favre R, et al. [Performance and indications of noninvasive prenatal testing using cell free circulating fetal DNA (cffDNA) for the detection of fetal trisomy 21, 18 and 13 in France]. *J Gynecol Obstet Biol Reprod (Paris)*. 2016;45:633–40.
30. Nshimyumukiza L, Menon S, Hina H, Rousseau F, Reinharz D. Cell-free DNA noninvasive prenatal screening for aneuploidy versus conventional screening: A systematic review of economic evaluations. *Clin Genet [Internet]*. [cited 2018 May 3];0. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1111/cge.13155>
31. Nshimyumukiza L, Beaumont J-A, Duplantie J, Langlois S, Little J, Audibert F, et al. Cell-Free DNA-Based Non-invasive Prenatal Screening for Common Aneuploidies in a Canadian Province: A Cost-Effectiveness Analysis. *J Obstet Gynaecol Can JOGC J Obstet Gynecol Can JOGC*. 2018;40:48–60.
32. Kostenko E, Chantraine F, Vandeweyer K, Schmid M, Lefevre A, Hertz D, et al. Clinical and Economic Impact of Adopting Noninvasive Prenatal Testing as a Primary Screening Method for Fetal Aneuploidies in the General Pregnancy Population. *Fetal Diagn Ther*. 2019;45:413–23.
33. John NM, Wright SJ, Gavan SP, Vass CM. The role of information provision in economic evaluations of non-invasive prenatal testing: a systematic review. *Eur J Health Econ HEPAC Health Econ Prev Care*. 2019;
34. Grosse SD, Wordsworth S, Payne K. Economic methods for valuing the outcomes of genetic testing: beyond cost-effectiveness analysis. *Genet Med Off J Am Coll Med Genet*. 2008;10:648–54.

Table 1: Values estimated for the variables included in the decision model to determine the most relevant threshold by performing cfDNA as the second-tier test

	Value (%)	Variations (%) [95%CI]	Costs (€)	References
Participation rate				
Karyotype	95.0	[90.0; 100.0]		Experts PEPRADE assumption
Sensitivity & Specificity				
Failure of cfDNA test, all causes combined	2.9	[0.8; 4.6]		Cuckle (2013)
FF < 4% (white, 60 kg, CRL 65 mm)	0.7	[0.2; 77.4]		Ashoor (2013)
Sensitivity of cfDNA test, all studies combined	99.2	[98.5; 99.6]		Gil (2015)
Specificity of cfDNA test, all studies combined	99.9	[99.87; 99.95]		Gil (2015)
Foetal loss after amniocentesis	0.11	[0.00; 0.26]		Akolekar (2015)
Foetal loss after trophoblast biopsy	0.22	[0.00; 1.16]		Akolekar (2015)
Costs and variations of costs included in the economic analysis				
cfDNA test			330	TNCB 2018
Amniocentesis + karyotyping			491	CCAM 2018; TNCB 2018
Chorionic villus sampling + karyotyping			593	CCAM 2018; TNCB 2018

cfDNA test: cell-free DNA, FF: foetal fraction, CRL: crown-rump length, CCAM: Classification Commune des Actes Médicaux (French classification of medical procedures), TNCB: Table Nationale de Codage de Biologie (National Table of Laboratory Codes)

Table 2: Performance of a conditional Down syndrome screening strategy including cell-free DNA testing as a second-tier test for first-trimester screening

Threshold value	n^a	cfDNA test	TP cfDNA test	FP cfDNA test	FN cfDNA test	Karyotype	Foetal loss	Total Down syndrome diagnosed
$\geq 1/50$	420	-	-	-	-	76	-	76
1/51-1/100	878	842	72	1	0	153	0	93
1/51-1/150	1430	1365	84	2	0	198	0	104
1/51-1/200	1992	1894	92	3	0	246	0	112
1/51-1/250	2627	2494	98	3	0	290	1	119
1/51-1/300	3108	2953	100	3	0	320	1	122
1/51-1/350	3668	3482	101	3	0	359	1	122
1/51-1/400	4300	4075	103	4	0	406	1	123
1/51-1/450	4905	4649	107	5	0	446	1	126
1/51-1/500	5480	5219	109	5	0	453	1	127
1/51-1/550	6049	5741	110	7	0	511	1	127
1/51-1/600	6698	6361	112	8	0	547	1	128

1/51-1/650	7274	6914	112	8	0	574	1	128
1/51-1/700	7863	7474	113	9	0	605	1	128
1/51-1/750	8429	8006	114	9	0	646	1	129
1/51-1/800	9029	8574	114	9	0	682	1	129
1/51-1/850	9611	9134	114	9	0	712	1	129
1/51-1/900	10187	9682	114	9	0	744	1	129
1/51-1/950	10754	10225	114	9	0	774	1	129
1/51-1/1000	11299	10744	114	9	0	803	1	129

cfDNA test: cell-free DNA test, TP: true positive, FP: false-positive, FN: false negative

^an: number of women concerned based on actual data in the simulation model

Table 3: Costs of the Down syndrome screening strategy considering different costs for cell-free DNA, CER and incremental cost.

Threshold value	cfDNA	cfDNA	cfDNA	cfDNA	CER	Cost per incremental DS diagnosed	
	cost 330	cost 50 euros ^a	cost 100 euros [†]	cost 200 euros ^a	cfDNA cost 330 euros ^a		
	euros [†]	(euros)	(euros)	(euros)	(euros)		
	(euros)						
1/51-1/100	359 900	124 140	166 240	250 440	21 171		
1/51-1/150	555 900	173 700	241 950	378 450	19 854		
1/51-1/200	756 585	226 265	320 965	510 365	21 016		
1/51-1/250	978 634	280 314	405 014	654 414	22 759		
1/51-1/300	1 141 171	314 331	461 981	757 281	24 808		
1/51-1/350	1 342 734	367 774	541 874	890 074	29 190		
1/51-1/400	1 564 459	423 459	627 209	1 034 709	33 286		
1/51-1/450	1 774 940	473 220	705 670	1 170 570	35 499		
1/51-1/500	1 966 576	505 256	766 206	1 288 106	38 560		191 637
1/51-1/550	2 171 089	563 609	850 659	1 424 759	42 570		396 149
1/51-1/600	2 395 506	614 426	932 476	1 568 576	46 067		310 283

1/51-1/650	2 592 166	656 246	1 001 946	1 693 346	49 849	408 613
1/51-1/700	2 793 200	700 480	1 074 180	1 821 580	53 715	509 130
1/51-1/750	2 992 982	751 302	1 151 602	1 952 202	56 471	406 014
1/51-1/800	3 197 876	797 156	1 225 856	2 083 256	60 337	474 312
1/51-1/850	3 398 009	840 489	1 297 189	2 210 589	64 113	541 023
1/51-1/900	3 595 882	884 922	1 369 022	2 337 222	67 847	606 981
1/51-1/950	3 793 281	930 281	1 441 531	2 464 031	71 571	672 781
1/51-1/1000	3 980 215	971 895	1 509 095	2 583 495	75 098	7352

cfDNA: cell-free DNA test, CER: cost-effectiveness ratio, DS: Down syndrome

^aThe simulation model used to calculate total cost at each threshold included the failure rate of the cfDNA test, which was performed a second time when the first failed.

Figure legends

Figure 1: Decision tree model of Down syndrome screening using cell-free DNA (cfDNA)

testing as a second-tier if first-trimester combined screening is positive, ranging from $\geq 1/50$

to $\geq 1/1000$ by step of $1/50$.

Figure 2: Cost of Down syndrome screening when performing cell-free DNA testing as a contingent strategy for a threshold of first-trimester combined screening 1/51-1/100 to 1/51-1/1000